

Math 222b Lecture Notes

Partial Differential Equations II

Professor: Maciej Zworski, Spring 2021

Scribe: Vishal Raman

Contents

1	January 19th, 2021	5
1.1	Review of Sobolev Spaces	5
1.2	Fourier Transform	6
2	December 21st, 2021	7
2.1	Plancherel's Theorem	7
2.2	Fourier Characterization of H^k spaces	7
3	January 26th, 2021	10
3.1	Sobolev Spaces, continued	10
3.2	Gagliardo-Nirenberg-Sobolev(GNS) Inequalities	10
3.3	Compactness	11
4	January 28th, 2021	12
4.1	Compactness	12
4.2	Rellich-Kondrachov	12
5	February 2nd, 2021	14
5.1	Rellich-Kondrachov, continued	14
5.2	Morrey's Inequality	15
6	February 4th, 2021	17
6.1	Morrey's Inequality, continued	17
7	February 9th, 2021	19
7.1	Fourier Transform proof of Morrey's Inequality	19
8	February 11th, 2021	21
8.1	Finishing Morrey's Inequality	21
8.2	Final Comments about Sobolev Spaces	21
8.3	Duality	22
8.4	Duality on Bounded Domains	22
9	February 16th, 2021	24
9.1	Calculus of Variations: Minimizing Distance in the Plane	24
9.2	Calculus of Variations: Minimizing Area in \mathbb{R}^3	24
9.3	Calculus of Variations: General Setup	25
9.4	Existence of Minimizers	26
10	February 18th, 2021	27
10.1	Second-Derivative Test	27
10.2	Convexity	28
10.3	Existence of Minimizers	28
11	February 23rd, 2021	29
11.1	Weak Convergence	29
11.2	Calculus of Variations	30
11.3	Getting around Lower Semicontinuity	30

12 February 25th, 2021	32
12.1 Existence of Minimizers	32
12.2 Uniqueness of Minimizers	32
12.3 The Euler-Lagrange Equation	33
13 March 2nd, 2021	35
13.1 Euler-Lagrange Equation, continued	35
14 March 4th, 2021	37
14.1 Regularity	37
14.2 Interior Regularity	37
15 March 9th, 2021	39
15.1 Inner Regularity	39
15.2 Higher Regularity	40
16 March 11th, 2021	41
16.1 Oscillatory Integrals	41
16.2 General Theory: Amplitudes	42
17 March 16th, 2021	43
17.1 Amplitudes of Oscillatory Integrals	43
17.2 A Cool Example	43
17.3 Topology of the Symbol Space	44
18 March 18th, 2021	45
18.1 Topology of the Symbol Space, continued	45
18.2 Phase Functions	45
19 March 30th, 2021	47
19.1 Oscillatory Integrals as Distributions	47
19.2 Some Remarks	48
20 April 1st, 2021	49
20.1 Oscillatory Integrals, continued	49
20.2 Fourier Integral Operators and Pseudodifferential Operators	50
21 April 6th, 2021	51
21.1 Operators on Oscillatory Integrals	51
21.2 Method of Stationary Phase	52
21.3 Steepest Descent	52
22 April 8th, 2021	54
22.1 Steepest Descent	54
22.2 Stirling's Formula	54
23 April 13th, 2021	56
23.1 Stationary Phase	56
23.2 Stationary Phase in Higher Dimensions	57
24 April 18th, 2021	58
24.1 Morse Lemma	58

24.2 Higher Dimensional Method of Stationary Phase	58
24.3 General Schrodinger Operators	59
25 April 20th, 2021	60
25.1 Pseudodifferential Operators	60

1 January 19th, 2021

1.1 Review of Sobolev Spaces

Definition 1.1. Given $u \in \mathcal{D}'(U)$ for $U \subseteq \mathbb{R}^n$ open: that means that $u : C_c^\infty(U) \rightarrow \mathbb{C}$ and for every compact set $K \subset\subset U$, $\exists C, N$ for all $\varphi \in C_0^\infty(K)$ such that

$$|u(\varphi)| \leq C \sup_{|\alpha| \leq N} |\partial^\alpha \varphi|.$$

Examples:

- Take $U = (0, 1)$ and take $u = \sum_{\mathbb{N}} \delta_{1/n}$, where $\delta_{1/n}(\varphi) = \varphi(1/n)$.
- Take $u \in L_{loc}^1(U)$, where $u(\varphi) = \int u \varphi$. Differentiation is defined formally though integration by parts as $\partial^\alpha u(\varphi) = (-1)^{|\alpha|} u(\partial^\alpha \varphi)$.

Definition 1.2. The Sobolev spaces $W^{k,p}(U) = \{u \in L_{loc}^1(U) : \partial^\alpha u \in L^p(U), \forall |\alpha| \leq k\}$, for $k \in \mathbb{N}_0$, $1 \leq p \leq \infty$. Note that differentiation is in the sense of distributions. We write $H^k(U) = W^{k,2}(U)$, which are Hilbert spaces with the inner product

$$\langle u, v \rangle = \sum_{|\alpha| \leq k} \int_U \partial^\alpha u \overline{\partial^\alpha v}.$$

Definition 1.3. $W_0^{k,p}(U) = \overline{C_c^\infty(U)}$, where the closure is with respect to the $W^{k,p}$ norm.

Theorem 1 (Approximation)

For $U \subset\subset \mathbb{R}^n$,

$$\overline{C^\infty(U) \cap W^{k,p}(U)} = W^{k,p}(U)$$

where the closure is with respect to the $W^{k,p}$.

If $\partial U \in C^1$, then we can improve up to

$$\overline{C^\infty(\overline{U}) \cap W^{k,p}(U)} = W^{k,p}(U)$$

Theorem 2 (Extension)

If $U \subset\subset \mathbb{R}^n$ and $\partial U \in C^1$, for $U \subset\subset V \subset\subset \mathbb{R}^n$, there exists $E : W^{1,p}(U) \rightarrow W^{1,p}(\mathbb{R}^n)$ such that $Eu|_U = u$ and the supp $u \subset\subset V$.

We can extend this to $W^{k,p}$ if the boundary is C^k .

Theorem 3 (Traces)

For $U \subset\subset \mathbb{R}^n$ with $\partial U \in C^1$, there exists $T : W^{1,p}(U) \rightarrow L^p(\partial U)$ which is linear and boundary such that for $u \in C(\overline{U}) \cap W^{1,p}$ $Tu = u|_{\partial U}$.

Example 1.4

For $U \subset\subset \mathbb{R}^n$, ∂U bounded,

$$H_0^1(U) = \{u \in H^1 : Tu = 0 \in L^2(\partial U)\}.$$

The converse of showing $Tu = 0$ implies H_0^1 is the more difficult one.

1.2 Fourier Transform

We first review the Fourier Transform. We define the Schwartz space:

$$\mathcal{S} = \{\varphi \in C^\infty(\mathbb{R}^n) : x^\alpha \partial^\beta \varphi \in L^\infty \forall \alpha, \beta \in \mathbb{N}^n\}.$$

For $\varphi \in \mathcal{S}$, we define

$$\widehat{\varphi}(\xi) = \int \varphi(x) e^{-ix \cdot \xi} dx.$$

Note that \mathcal{F} , the Fourier transform is invertible on \mathcal{S} . The key properties of the Fourier transform are

$$\mathcal{F}(1/i\partial x \varphi) = \xi \mathcal{F}\varphi, \mathcal{F}(x\varphi) = -1/i\partial_\xi \mathcal{F}\varphi.$$

We also have

$$\mathcal{F}^{-1} = \frac{R\mathcal{F}}{(2\pi)^n}, R\varphi(x) = \varphi(-x).$$

We define \mathcal{S}' onto \mathbb{C} so that for $u \in \mathcal{S}'$, there exists C, N such that

$$|u(\varphi)| \leq C \sup_{|\alpha|, |\beta| \leq N} |x^\alpha \partial^\beta \varphi|.$$

Note that $\mathcal{S}' \subset \mathcal{D}'$.

Definition 1.5. $\mathcal{F} : \mathcal{S}' \rightarrow \mathcal{S}'$ by $\widehat{u}(\varphi) = u(\widehat{\varphi})$.

Examples:

- $\widehat{\delta}_0(\varphi) = \delta_0(\widehat{\varphi}) = \widehat{\varphi}(0) = \int \varphi = 1(\varphi)$.
- Take \mathbb{R}^2 and consider $u(x) = \frac{1}{|x|}$. This function is in L^1_{loc} . If we multiply by $(1 + |x|)^{-2}u \in L^1(\mathbb{R}^n)$, it follows that $u \in \mathcal{S}'$, since

$$|u(\varphi)| = \left| \int (1 + |x|)^{-2}u(1 + |u|)^2\varphi \right| \leq C \sup(1 + |x|)^2\varphi.$$

Now, we compute $\widehat{u} \in \mathcal{S}'$. Since \mathcal{F} is continuous on \mathcal{S}' , we approximate u and hope the result converges to the desired result. Define $u_\epsilon \rightarrow u$ in \mathcal{S}' for $u_\epsilon \in L^1$.

Try $u_\epsilon(x) = \frac{e^{-\epsilon|x|^2/2}}{|x|} \in L^1$ for $\epsilon > 0$. We want to calculate \widehat{u}_ϵ and take the limit as $\epsilon \rightarrow 0^+$. We can evaluate the integral by converting to polar coordinates and completing the square. Unfortunately, it reduces to an integral that is too hard, but we will learn asymptotics of the integral as $\epsilon \rightarrow 0$. We find that $\widehat{u}(\xi) = 2\pi/|\xi|$.

We can approach this differently. Note that $u = 1/|x|$ is homogeneous: $u(tx) = t^a u(x)$ for $t > 0$, for functions. For distributions, we have that for $\varphi \in \mathcal{S}$, $u(\varphi(\cdot/t)t^{-n}) = t^a u(\varphi)$ for $t > 0$. For the Fourier Transform, if $u \in \mathcal{S}'(\mathbb{R}^n)$ is homogeneous of degree a , then \widehat{u} is homogeneous of degree $-n - a$. It follows that our Fourier transform is of degree -1 .

Furthermore, note that $1/|x|$ is spherically symmetric, and the Fourier transform preserves spherical symmetry (note that the Jacobian factor for rotations is 1). It follows that the Fourier transform is also spherically symmetric. It follows that

$$\mathcal{F}(1/|x|) = C/|\xi| + \sum_{|\alpha| \leq N} c_\alpha \delta_0^{(\alpha)},$$

but delta terms have too much homogeneity.

2 December 21st, 2021

2.1 Plancherel's Theorem

Recall that the Fourier transform is an isomorphism on \mathcal{S} - it is a bounded linear operator whose inverse is also bounded.

Note that

$$\int \widehat{u}(\xi) \overline{\widehat{\varphi}(\xi)} d\xi = \iiint u(x) \overline{\varphi(y)} e^{-i(x-y)\xi} dx dy d\xi$$

In the sense of distributions, $\int e^{-i(x-y)\xi} d\xi = (2\pi)^n \delta(x-y)$. Hence,

$$\iiint u(x) \overline{\varphi(y)} e^{-i(x-y)\xi} dx dy d\xi = (2\pi)^n \int u(x) \overline{\varphi(x)} dx.$$

For $u, \varphi \in \mathcal{S}$, we have the following:

$$\langle \widehat{u}, \widehat{\varphi} \rangle = (2\pi)^n \langle u, \varphi \rangle.$$

This implies that

$$\|\widehat{u}\|_2 = (2\pi)^{n/2} \|u\|_2, u \in \mathcal{S}.$$

If $u_n \rightarrow u$ in L^2 then $u_n \rightarrow u$ in \mathcal{S}' by the Cauchy-Schwartz inequality. It follows that $\widehat{u}_n \rightarrow \widehat{u}$ in \mathcal{S}' but our formula shows that \widehat{u} is in L^2 . Hence, $\mathcal{F} : L^2 \rightarrow L^2$ and for $u, v \in L^2$, $\langle \widehat{u}, \widehat{v} \rangle = (2\pi)^n \langle u, v \rangle$.

Recall last time, we were finding the Fourier transform of $u(x) = 1/|x|$ in \mathbb{R}^2 . For $u \in \mathcal{S}'(\mathbb{R}^n)$ homogeneous of degree a , $\widehat{u} \in \mathcal{S}'(\mathbb{R}^n)$ is homogeneous of degree $-n-a$. In our example, It follows that $\widehat{u}(\xi)$ is homogeneous of degree -1 . We also observed that u is invariant under rotations so it follows that \widehat{u} is invariant under rotations.

A function is homogeneous of degree -1 if $v(k\theta) = \frac{a(\theta)}{r}$. Since our function is invariant under rotations, $\widehat{u}(\xi) = \frac{c}{|\xi|}$ away from zero. It follows from our previous argument that $\widehat{u}(\xi) = \frac{c}{|\xi|}$ since δ terms have homogeneity of at least -2 .

Note that $\langle u, \varphi \rangle = (2\pi)^2 \langle \widehat{u}, \widehat{\varphi} \rangle$ and we find \widehat{u} by choosing an appropriate φ .

$$\begin{aligned} \int_{\mathbb{R}^2} \frac{\varphi(x)}{|x|} dx &= \int_0^{2\pi} \int_0^\infty \varphi(r) dr d\theta \\ &= 2\pi \int_0^\infty \varphi(r) dr. \end{aligned}$$

Choosing $\varphi(r) = e^{-r^2/2}$, we find that the integral is $(2\pi)^{3/2}$.

Evaluating the other side,

$$\widehat{\varphi}(\xi) = \int_{\mathbb{R}^2} e^{-|x|^2/2 - ix \cdot \xi} dx = \int e^{-\frac{1}{2}(x+i\xi)^2 - \frac{1}{2}|\xi|^2} = 2\pi \int e^{-|\xi|^2/2} = (2\pi)^{5/2}.$$

It follows that $c = 2\pi$.

2.2 Fourier Characterization of H^k spaces

Theorem 4

$$H^k(\mathbb{R}^n) = \{u \in \mathcal{S}'(\mathbb{R}^n) : (1 + |\xi|^2)^{k/2} \widehat{u} \in L^2\}.$$

Proof. Suppose that $\partial^\alpha u \in L^2$ for $|\alpha| \leq k$. We know that $\|u\|_2 = (2\pi)^{-n/2} \|\widehat{u}\|$. It follows that $\widehat{\partial^\alpha u} \in L^2$. Note that $\widehat{\partial^\alpha u} = i^{|\alpha|} x^{i\alpha} \widehat{u} \in L^2$ for all $|\alpha| \leq k$.

Hence,

$$(1 + |\xi|^2)^{k/2} \leq C_{n,k} \sup_{|\alpha| \leq k} |\xi^\alpha|.$$

So it follows that $(1 + |\xi|^2)^{k/2} \widehat{u} \in L^2$.

Now, suppose $(1 + |\xi|^2)^{k/2} \widehat{u} \in L^2$. It follows that $|\xi^\alpha| \leq C_{k,\alpha} (1 + |\xi|^2)^{k/2}$ for $|\alpha| \leq k$. Hence $\xi^\alpha \widehat{u} \in L^2$ so it follows that $\partial^\alpha u \in L^2$ by Plancherel's Theorem. \square

Remark 2.1. We use the notation $\langle \xi \rangle = (1 + |\xi|^2)^{1/2}$.

Note that the definition does not require $k \in \mathbb{N}$.

Definition 2.2. $H^s(\mathbb{R}^n) = \{u \in \mathcal{S}' : \langle \xi \rangle^s \widehat{u} \in L^2\}$, $s \in \mathbb{R}$.

Theorem 5

Suppose $u \in H^s(\mathbb{R}^n)$ and $s > \frac{1}{2}$. Then $v(y) = u(0, y)$, $y \in \mathbb{R}^{n-1}$ satisfies $v \in H^{s-1/2}(\mathbb{R}^{n-1})$.

Remark 2.3. We should define $Tu(y) = u(0, y)$ for $u \in \mathcal{S}$. Then $T : H^s(\mathbb{R}^n) \rightarrow H^{s-1/2}(\mathbb{R}^{n-1})$ if $s > 1/2$.

Proof. Take $u \in \mathcal{S}$. We wish to show that $\|v\|_{H^{s-1/2}(\mathbb{R}^{n-1})} \leq C \|u\|_{H^s(\mathbb{R}^n)}$.

Note that

$$\widehat{v}(\eta) = \int_{\mathbb{R}^{n-1}} u(0, y) e^{-iy \cdot \eta} dy$$

and by the Fourier Inversion Formula

$$u(0, y) = (2\pi)^{-n} \int_{\mathbb{R}^n} \widehat{u}(\xi_1, \xi') e^{iy \cdot \xi'} d\xi_1 d\xi',$$

so it follows that

$$\begin{aligned} \widehat{v}(\eta) &= (2\pi)^{-n} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} \widehat{u}(\xi_1, \xi') e^{-iy \cdot (\eta - \xi')} d\xi dy \\ &= (2\pi)^{-n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^{n-1}} \widehat{u}(\xi_1, \xi') e^{iy \cdot (\xi' - \eta)} dy d\xi \\ &= (2\pi)^{-1} \int_{\mathbb{R}^n} \widehat{u}(\xi_1, \xi') \delta_{\xi' = \eta} d\xi \\ &= (2\pi)^{-1} \int_{\mathbb{R}} \widehat{u}(\xi_1, \eta) d\xi_1. \end{aligned}$$

Note that up to constants

$$\|v\|_{H^{s-1/2}}^2 = \int_{\mathbb{R}^{n-1}} \langle \eta \rangle^{2s-1} |\widehat{v}(\eta)|^2 d\eta = \int_{\mathbb{R}^{n-1}} \langle \eta \rangle^{2s-1} \left| \int_{\mathbb{R}} \widehat{u}(\xi_1, \eta) d\xi_1 \right|^2 d\eta.$$

Then,

$$\begin{aligned}
& \int_{\mathbb{R}^{n-1}} \langle \eta \rangle^{2s-1} \left| \int \widehat{u}(\xi_1, \eta) d\xi_1 \right|^2 d\eta \\
&= \int \langle \eta \rangle^{2s-1} \left| \int \widehat{u}(\xi, \eta) (1 + |\xi_1|^2 + |\eta|^2)^{s/2} (1 + |\xi_1|^2 + |\eta|^2)^{-s/2} d\xi_1 \right|^2 d\eta \\
&\leq \int \langle \eta \rangle^{2s-1} \int |\widehat{u}(\xi_1, \eta)|^2 (1 + |\xi_1|^2 + |\eta|^2)^s d\xi_1 \int (1 + |\xi_1|^2 + |\eta|^2)^{-s} d\xi_1 d\eta \\
&\leq \int \langle \eta \rangle^{2s-1} \langle \eta \rangle^{-2s+1} \int |\widehat{u}(\xi_1, \eta)|^2 (1 + |\xi_1|^2 + |\eta|^2)^s d\xi_1 \int (1 + u^2)^{-s} dud\eta \\
&= \int |\widehat{u}(\xi)|^2 \langle \xi \rangle^{2s} d\xi = \|u\|_{H^s}^2,
\end{aligned}$$

since

$$\int |\widehat{u}(\xi_1, \eta)|^2 (1 + |\xi_1|^2 + |\eta|^2)^s d\xi_1 d\eta = \int |\widehat{u}(\xi)|^2 \langle \xi \rangle^{2s} d\xi.$$

□

3 January 26th, 2021

3.1 Sobolev Spaces, continued

Recall, we have $U \subset \mathbb{R}^n$ open. We typically assume U is bounded and $\partial U \in C^1$. For these spaces, we define

$$W^{k,p}(U) = \{u \in \mathcal{D}' : \partial^\alpha u \in L^p(U), |\alpha| \leq k\}.$$

Recall the extension property: there exists a map $E : W^{1,p}(U) \rightarrow W^{1,p}(\mathbb{R}^n)$ such that $Eu|_U = u$ and $u = 0$ for $|x| > R$ for some R with $U \subset\subset B(0, R)$.

We also consider the $H^s(\mathbb{R}^n)$, the fractional Sobolev spaces: $\{u \in \mathcal{S}'(\mathbb{R}^n) : \langle \xi \rangle^s \widehat{u} \in L^2\}$. This is a Hilbert space with the norm

$$\|u\|_{H^s}^2 = \int \langle \xi \rangle^{2s} |\widehat{u}(\xi)|^2 d\xi.$$

Last time, we showed that If we have $u \in H^s(\mathbb{R}^n)$ and $s > 1/2$, then $v(y) : u(0, y), y \in \mathbb{R}^{n-1}$ satisfies $v \in H^{s-1/2}(\mathbb{R}^{n-1})$. Today, we will show that $H^s(\mathbb{R}^n) \subset C_0(\mathbb{R}^n)$ if $s > n/2$, where C_0 denotes continuous functions vanishing at infinity. This means that there exists $T : H^s(\mathbb{R}^n) \rightarrow H^{s-1/2}(\mathbb{R}^{n-1})$ such that for $u \in \mathcal{S}$, $Tu(y) = u(0, y)$.

Theorem 6

$H^s(\mathbb{R}^n) \subset C_0(\mathbb{R}^n)$ if $s > n/2$.

Proof. We first prove that if $\langle \xi \rangle^s \widehat{u} \in L^2, s > n/2$ then $\widehat{u} \in L^1(\mathbb{R}^n)$.

$$\int_{\mathbb{R}^n} |\widehat{u}| d\xi = \int_{\mathbb{R}^n} \langle \xi \rangle^{-s} \langle \xi \rangle^s |\widehat{u}| d\xi \leq \|\langle \xi \rangle^{-s}\|_2 \|u\|_{H^s}.$$

The first term is finish precisely when $s > n/2$ [exercise: convert to polar coordinates]. This implies that $u \in L^\infty(\mathbb{R}^n)$, following from the Fourier Inversion formula.

We know that $x \mapsto \widehat{u}(\xi)e^{ix\xi}$ is continuous so it follows that $x \mapsto u(x)$ is continuous by the dominated convergence theorem. Finally $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$ by the Riemann-Lebesgue lemma: if $\widehat{u} \in L^1(\mathbb{R}^n)$, then $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$.

Proof. Recall $\mathcal{S}(\mathbb{R}^n) \subset L^1(\mathbb{R}^n)$ is dense. Taking $v \in L^1$, taking $v_R = v(x)1_{B(0,R)}(x)$. Then $v_R \rightarrow v$ y the dominated convergence theorem. Now take $\varphi \in C_c^\infty$ with $\varphi \geq 0, \int \varphi = 1$ wth $\varphi_\epsilon(x) = \frac{1}{\epsilon^n} \varphi(x/\epsilon)$. Taking $v_{R,\epsilon} = v_R * \varphi_\epsilon \in C_c^\infty(\mathbb{R}^n)$ and $v_R * \varphi_\epsilon \rightarrow v_R$ in L^1 as $\epsilon \rightarrow 0$.

Hence, we can take $v \in \mathcal{S}$ so that $\|\widehat{v} - \widehat{u}\|_{L^1} < \epsilon/2$. Now, $|v(x)| < \epsilon/2$ if $|x| > R$, hence

$$|u(x)| \leq |u(x) - v(x)| + |v(x)| < C\epsilon + \epsilon/2$$

which goes to 0 as we send $\epsilon \rightarrow 0$. □

□

3.2 Gagliardo-Nirenberg-Sobolev(GNS) Inequalities

Theorem 3.1

If $1 \leq p < n$ and we define $p^* = \frac{np}{n-p}$, then there exists $C = C(p, n)$ so that for all $u \in C_c^\infty(\mathbb{R}^n)$,

$$\|u\|_{L^{p^*}} \leq C \|\nabla u\|_p.$$

Remark 3.2. We can find the value of p^* without doing the computation through scaling. Take $u_\lambda(x) = u(\lambda x)$. We have that $\|u_\lambda\|_{p^*} \leq C \|\nabla(u_\lambda)\|_p$. Then, evaluate both sides and compare the exponent on λ .

Note that the result is not true for $p = n > 1$. It is true for $p = n = 1$.

Theorem 3.3 (Morrey's Inequality)

For $n < p \leq \infty$, there exists $C = C(p, n)$ such that for $u \in C^1(\mathbb{R}^n)$, we have

$$\|u\|_{C^\gamma(\mathbb{R}^n)} \leq C(\|u\|_p + \|\nabla u\|_p),$$

where $\gamma = 1 - \frac{n}{p}$, where

$$\|u\|_{C^\gamma(\mathbb{R}^n)} = \sup |u| + \sup_{x \neq y} \frac{|u(x) - u(y)|}{|x - y|^\gamma}.$$

Theorem 7 (General Formulation)

Take $U \subset \subset \mathbb{R}^n$ with $\partial U \in C^1$. Take $n \in W^{k,p}(U)$.

- if $k < n/p$, then $u \in L^q(U)$ where $1/q \geq 1/p - k/n$ and $\|u\|_{L^q(U)} \leq C\|u\|_{W^{k,p}}$.
- $k > n/p$, then $u \in C^{k-[n/p]-1, \gamma}(\bar{U})$ where $\gamma = [n/p] + 1 - n/p$ if $n/p \notin \mathbb{N}$ and $1 - \delta$ for all $\delta > 0$ if $n/p \in \mathbb{N}$.

3.3 Compactness

Definition 3.4. Let B be a Banach space. A subset $K \subset B$ is compact if for every sequence $\{u_n\} \subset K$ such that $\|u_n\|_B \leq C$, there exists a convergence subsequence $u_{n_k} \rightarrow u \in B$.

Remark 3.5. If $\{u : \|u\|_B \leq 1\} \subset B$ is compact, then B is finite dimensional. We can have a space $B' \subset B$ and $\{u \in B' : \|u\|_{B'} \leq 1\}$ compact in B . If we have a sequence $\{u_n\} \subset B'$ and $\|u_n\|_{B'} \leq C$ then there exists $n_k, u \in B$ such that $\|u_{n_k} - u\|_B \rightarrow 0$.

We will take $B = L^q(U)$ where $1 \leq q < p^*$ and $B' = W^{1,p}(U)$.

Theorem 8 (Rellich-Kondrachov)

The unit ball in $W^{1,p}(U)$ is compact in $L^q(U)$ for **bounded** U .

4 January 28th, 2021

Recall the GNS inequality: if $1 \leq p < n$, $p^* = \frac{np}{n-p}$, there exists $C = C(n, p)$ for all $u \in C_c^\infty(\mathbb{R}^n)$ so that $\|u\|_{L^{p^*}} \leq C \|\nabla u\|_{L^p}$.

If $U \Subset \mathbb{R}^n$, $\partial U \in C^1$, then there exists $C = C(n, p, U)$ such that $L^q(U) \supset W^{1,p}(U)$ for $1 \leq q \leq p^*$.

4.1 Compactness

Suppose B is a Banach space and $B' \subset B$ another Banach space. We say that the inclusion $B' \subset B$ is compact if bounded sets in B' are precompact in B . In other words, for a sequence $\{u_n\} \subset B'$ with $\|u_n\|_{B'} \leq M$, there exists a subsequence u_{n_k} and $u \in B$ such that $u_{n_k} \rightarrow u$ in B .

Example 4.1

Take $B = C([-1, 1])$, $B' = C^1([-1, 1])$ with the supremum norm on B and $\|u\|_{B'} = \sup_{|x| \leq 1} (|u(x)| + |u'(x)|)$.

The inclusion is compact: if we have $\|u_n\|_{B'} \leq C$, by the mean value theorem, $|u_n(x)| \leq C$ and $|u_n(x) - u_n(y)| \leq C|x - y|$. By Arzela-Ascoli, there exists a subsequence u_{n_k} and $u \in C$ so that $\|u_{n_k} - u\|_{C([-1, 1])} \rightarrow 0$.

Example 4.2

In the previous example, take $u_n(x) = |x|1_{|x| > 1/n} + (\frac{nx^2}{2} + \frac{1}{n})1_{|x| \leq 1/n}$.

Then $u_n \in C^1[-1, 1]$ and $\|u_n\|_{C^1[-1, 1]} \leq 2$. We can take a subsequence $n_k = k$ and $u(x) = |x| \in C[-1, 1] \setminus C^1[-1, 1]$ where $u_{n_k} \rightarrow u \in C$.

Given a Banach space B , we have the dual space $B^* = \{\text{linear } u : B \rightarrow \mathbb{C} \mid \forall x \in B, |u(x)| \leq C\|x\|_B\}$. This is also a Banach space.

Theorem 9 (Banach-Alaoglu)

Suppose $\|u_n\|_{B^*} \leq M$. Then, there exists a subsequence u_{n_k} and $u \in B^*$ such that for all $x \in B$, $u_{n_k}(x) \rightarrow u(x)$.

4.2 Rellich-Kondrachov

Theorem 10 (Rellich-Kondrachov)

If $U \Subset \mathbb{R}^n$, $\partial U \in C^1$, then for $1 \leq q < p^*$, $L^q(U) \supset W^{1,p}(U)$ is a compact inclusion.

Proof. Take $p = 2$. Then $p^* = \frac{2n}{n-2} > 2$. First, suppose $\bar{U} \Subset B(0, R)$. We can assume $R = 1$. Suppose we have a sequence $\|v_n\|_{H^1(U)} \leq 1$. There exists a sequence $u_n \in H^1(\mathbb{R}^n)$ such that $u_n|_U = v_n$, $\|u_n\|_{H^1(\mathbb{R}^n)} \leq 1$ and $\text{supp } u_n \subset B(0, 1)$ (this is the extension operator).

We have $u_n \in H^1(\mathbb{R}^n)$, $\|u_n\|_{H^1} \leq 1$, $\text{supp } u_n \subset B(0, R)$. We want n_k , $u \in L^2(\mathbb{R}^n)$ such that $u_{n_k} \rightarrow u$ in L^2 . We claim that $u(x) = (2\pi)^{-n} \sum_{m \in \mathbb{Z}^n} \hat{u}(m) e^{im \cdot x}$ for $x \in B(0, 1) \subset [-\pi, \pi]^n$ with convergence in \mathcal{D}' .

Alternatively,

$$\begin{aligned} \int u(x)\overline{\varphi(x)} dx &= (2\pi)^{-n} \sum_{m \in \mathbb{Z}^n} \widehat{u}(m) \int \overline{\varphi(x)} e^{im \cdot x} dx \\ &= (2\pi)^{-n} \sum_{m \in \mathbb{Z}^n} \widehat{u}(m) \widehat{\overline{\varphi}}(m). \end{aligned}$$

For $u, v \in L^2$, $\int u(x)\overline{v(x)} dx = (2\pi)^{-n} \sum_{m \in \mathbb{Z}^n} \widehat{u}(m) \overline{\widehat{v}(m)}$.

Recall the Poisson summation formula: for $n = 1, a \neq 0$,

$$\sum_{k \in \mathbb{Z}} e^{ikax} = \frac{2\pi}{a} \sum_{k \in \mathbb{Z}} \delta(x - 2\pi k/a)$$

in the distributional sense.

Note that $(1 - e^{iax}) \sum_k e^{ikax} = \sum_k e^{ikax} - \sum_k e^{i(k+1)ax} = 0$. We can rewrite this as $-2ie^{-iax/2} \sin(ax/2) \sum_k e^{ikax} = 0$. Let $w(x) = \sum_k e^{ikax}$, so it follows that $\text{supp } w \subset \{\frac{2\pi}{a}k\}_{k \in \mathbb{Z}}$. It follows that $w(x)$ is the sum of delta functions supported at $2\pi k/a$ for $k \in \mathbb{Z}$ up to constants.

Furthermore, note that $w(x + 2\pi a) = w(x)$. So it follows that the constants are independent of the index. To find the constant, for some function, replace $\varphi(\cdot)$ with $\varphi(\cdot + x)$. Then the right side is $c \sum_{k \in \mathbb{Z}} \varphi(2\pi k/a + x)$. Note that $\widehat{\varphi}(\cdot + x)(\xi) = e^{ix\xi} \widehat{\varphi}(\xi)$. It follows that the left hand side is $\sum_{k \in \mathbb{Z}} \widehat{\varphi}(ka) e^{ikax}$. Now suppose $\text{supp } \varphi \in C_c^\infty((0, 2\pi/a))$. Integrating both sides, the left side is $2\pi/a \widehat{\varphi}(0)$. The right side is $c \int \varphi(x) dx = c(a) \widehat{\varphi}(0)$. Thus, $c = \frac{2\pi}{a}$.

The Poisson summation formula is more generally $\sum_{k \in \mathbb{Z}^k} e^{iak \cdot x} = (2\pi/a)^n \sum_{k \in \mathbb{Z}^d} \delta(x - 2\pi a/k)$.

Applying this to a φ gives our desired claim from earlier. it follows that $\frac{1}{(2\pi)^n} \sum_{k \in \mathbb{Z}^n} \widehat{u}(k) \overline{\widehat{v}(k)} = \int u(x)\overline{v(x)} dx$ with $u, v \in L^2$, $\text{supp } u, v \in [-\pi, \pi]^n$.

Note that for $u \in L^2$, we have the Plancherel formula,

$$\|u\|_2^2 = \int |u(x)|^2 dx = \frac{1}{2\pi} \sum_{n \in \mathbb{Z}^n} |\widehat{u}(m)|^2.$$

For $u \in H^1(\mathbb{R}^n)$ and $\text{supp } u \subset B(0, 1)$, then $u \in L^2$ and $\partial^\alpha u \in L^2$, for all $|\alpha| = 1$,

$$\|\partial^\alpha u\|_{L^2}^2 = (2\pi)^n \sum_{m \in \mathbb{Z}^n} |\widehat{\partial^\alpha u}(m)|^2 = (2\pi)^{-n} \sum_{m \in \mathbb{Z}^n} |m^\alpha \widehat{u}(m)|^2.$$

Claim: Under these assumptions, $\|u\|_{H^1}^2 = \Theta(\sum_{m \in \mathbb{Z}^n} \langle m \rangle^2 |\widehat{u}(m)|^2)$ □

5 February 2nd, 2021

Recall the following:

- GNS inequality: For $U \in \mathbb{R}^n$, $\partial U \in C^1$, $1 \leq p < n$, $p^* = \frac{np}{n-p} > p$,

$$\|u_{-p^*} \leq C(\|u\|_p + \|\nabla u\|_p).$$

- R-K Theorem: For $1 \leq p < n$, $1 \leq q < p^*$,

$$W^{1,p}(U) \subset L^q(U)$$

is compact: If we have $\{u_n\} \subset W^{1,p}(U)$ and $\|u_n\|_{W^{1,p}} \leq C$, there exists a subsequence u_{n_k} , $u \in L^q$ such that $\|u_{n_k} - u\|_q \rightarrow 0$.

5.1 Rellich-Kondrachov, continued

Last time, we considered the special case of $\{u_n\} \subset H^1(\mathbb{R}^n)$ such that $\text{supp } u_n \subset B(0, R)$ and $\|u_n\|_{H^1} \leq C$, which implies that there exists a subsequence u_{n_k} and $u \in L^2(\mathbb{R}^n)$ such that $\|u_n - u\|_{L^2} \rightarrow 0$. We continue the proof of the special case.

Proof. Recall that we showed that if $u \in C_0^\infty((-\pi, \pi)^n)$, we can write $u(x) = (2\pi)^{-n} \sum \widehat{u}(n)e^{in \cdot x}$. Then

$$\int u(x)\overline{v(x)} dx = \frac{1}{(2\pi)^n} \sum \widehat{u}(n)\overline{\widehat{v}(n)}$$

and

$$\int |\nabla u(x)|^2 dx = \frac{1}{(2\pi)^n} \sum |n|^2 |\widehat{u}(n)|^2.$$

For $u \in H^1$ with $\text{supp } u \in B(0, 1)$,

$$\|u\|_{H^1}^2 = \frac{1}{(2\pi)^n} \sum \langle n \rangle^2 |\widehat{u}(n)|^2.$$

$$\|u_n\|_{H^1}^2 = \sum_{\ell \in \mathbb{Z}^n} \langle \ell \rangle^2 |\widehat{u}_n(\ell)|^2 \leq C.$$

$$\|v\|_{L^2}^2 = \sum_{\ell \in \mathbb{Z}^n} |\widehat{v}(\ell)|^2.$$

We want to show that there exists n_k such that $\|u_{n_k} - u_{n_p}\|_{L^2} \rightarrow 0$ as $k, p \rightarrow \infty$.

We introduce an operator $\Pi_p u(x) = (2\pi)^{-n} \sum_{|\ell| \leq p} \widehat{u}(\ell)e^{i\ell \cdot x}$. We can think of $\Pi_p : L^2([-\pi, \pi]^n) \rightarrow \mathbb{C}^{N_p}$. N_p can be found through combinatorial methods (left as an exercise) [should be $\binom{n+p}{p}$ or something like that].

We have the following estimate:

$$\|(I - \Pi_p)u\|_2 \leq \langle p \rangle^{-2} \|u\|_{H^1}^2.$$

This is because

$$(2\pi)^{-n} \sum_{|\ell| > p} |\widehat{u}(\ell)|^2 = (2\pi)^{-n} \sum_{|\ell| > p} \langle \ell \rangle^{-2} \langle \ell \rangle^2 |\widehat{u}(\ell)|^2 \leq \langle p \rangle^{-2} \|u\|_{H^1}^2.$$

Now, we find the Cauchy subsequence.

1. For all p , we have $\|\Pi_p u_n\|_{\mathbb{C}^{N_p}} \leq \|u_n\|_2 \leq \|u_n\|_{H^1} \leq C$. Then $\{|z| \leq C\} \subset \mathbb{C}^{N_p}$ is compact. It follows that we can choose subsequences $\{n_k^{p+1}\} \subset \{n_k^p\}$ such that $\Pi_p u_k^p$ converges and $\limsup_{k,\ell} \|u_k^p - u_\ell^p\| \leq C\langle p \rangle^{-2}$, which follows from the triangle inequality. [let $u_a^b = u_{n_a^b}$]
2. We choose $n_k = n_k^k$. It follows that $\limsup_{k,\ell \rightarrow \infty} \|u_{n_k} - u_{n_\ell}\|_2 = 0$, since

□

5.2 Morrey's Inequality

Theorem 11 (Morrey's Inequality)

Suppose $u \in L^p(\mathbb{R}^n)$, $\nabla u \in L^p(\mathbb{R}^n)$ and $n < p \leq \infty$. Then there exists $u^* \in C^{0,\gamma}(\mathbb{R}^n)$, with $\gamma = 1 - \frac{n}{p}$ such that $u = u^*$ almost everywhere and $\|u^*\|_{C^{0,\gamma}} \leq \|u\|_p + \|\nabla u\|_p$.

Remark 5.1. Recall

$$\|u\|_{C^{0,\gamma}} = \sup |u(x)| + \sup_{x \neq y} \frac{|u(x) - u(y)|}{|x - y|^\gamma}.$$

Proof. We use the Littlewood-Paley Decomposition.

Lemma 5.2 (Dyadic Partitions of Identity)

There exists a function $\psi_0 \in C_c^\infty(\mathbb{R})$, $\psi \in C_c^\infty(\mathbb{R} \setminus \{0\})$ such that

$$\psi_0(\xi) + \sum_{j=0}^{\infty} \psi(2^{-j}|\xi|) = 1.$$

Proof. Choose $\varphi_0 \in C_c^\infty((-1, 1))$ with $0 \leq \varphi_0 \leq 1$ and $\varphi_0(p) = 1$, $|p| \leq 1/2$.

Choose a new function

$$\varphi_1(p) = \sum_{j \in \mathbb{Z}} \varphi_0(p - j) \geq 1.$$

Note that $\varphi_1(p - k) = \varphi_1(p)$ for $k \in \mathbb{Z}$. Choose $\varphi(p) = \frac{\varphi_0(p)}{\varphi_1(p)}$.

Then

$$\sum_{j \in \mathbb{R}} \varphi(p - j) = \sum_{j \in \mathbb{R}} \frac{\varphi_0(p - j)}{\varphi_1(p - j)} = \frac{1}{\varphi_1(p)} \sum_{j \in \mathbb{R}} \varphi_0(p - j) = 1.$$

Define $\psi(r) = \varphi(\frac{\log r}{\log 2})$ for positive r . Notice that $\psi \in C_c^\infty((0, \infty))$. This gives that

$$\sum_{j \in \mathbb{Z}} \psi(2^{-j}r) = 1.$$

Define $\psi_0(r) = 1 - \sum_{j=0}^{\infty} \psi(2^{-j}r)$. Note that $\psi_0(r) = 1$ for $r < 1/2$ and $\psi_0(r) = 0$ for $r > 1$. It follows that ψ_0 and ψ satisfy the conditions. □

We can extend the Dyadic Partitions of Identity in \mathbb{R}^n in the natural way. We then define the Littlewood-Paley Decomposition as

$$u = \psi_0(D)u + \sum_{j=1}^{\infty} \psi(2^{-j}D)u$$

where for $a \in L^\infty(\mathbb{R}^n)$, $a(D)u = \mathcal{F}^{-1}(a(\xi)\widehat{u}(\xi))$ where $D_x = 1/i\partial_x$ and $\widehat{Du} = \xi\widehat{u}$. \square

6 February 4th, 2021

6.1 Morrey's Inequality, continued

Recall the statement of the theorem.

Theorem 12 (Morrey's Inequality)

Suppose $u \in L^p(\mathbb{R}^n)$, $\nabla u \in L^p(\mathbb{R}^n)$ and $n < p \leq \infty$. Then there exists $u^* \in C^{0,\gamma}(\mathbb{R}^n)$, with $\gamma = 1 - \frac{n}{p}$ such that $u = u^*$ almost everywhere and $\|u^*\|_{C^{0,\gamma}} \leq \|u\|_p + \|\nabla u\|_p$.

Proof. Recall for $u \in \mathcal{S}(\mathbb{R})$, $\widehat{D_{x_j} u}(\xi) = \xi_j \widehat{u}(\xi)$ where $D_{x_j} = \frac{1}{i} \partial_{x_j}$.

We define a **Fourier multiplier** $a \in L^\infty(\mathbb{R}^n)$ so that $a(D)u = \mathcal{F}^{-1}(a(\xi)\widehat{u}(\xi))$ for $u \in \mathcal{S}$. Note that for $a \in L^\infty$, $\|a(D)u\|_{L^2} \leq \sup |a| \|u\|_{L^2}$. For $\psi \in \mathcal{S}(\mathbb{R}^n)$, if we take $u \in \mathcal{S}'$, then $\psi(D)u \in \mathcal{S}'$, and $\psi(\xi)\widehat{u}(\xi) \in \mathcal{S}'$.

Recall the Littlewood - Paley Decomposition. We had a lemma: there exists $\psi_0 \in C_c^\infty(\mathbb{R})$ and $\psi \in C_c^\infty(\mathbb{R} \setminus \{0\})$ such that for all $\xi \in \mathbb{R}^n$,

$$\psi_0(|\xi|) + \sum_{j=0}^{\infty} \psi(2^{-j}|\xi|) = 1.$$

Slightly abusing notation, we will write $\psi_0(\xi) = \psi_0(|\xi|)$ and $\psi(\xi) = \psi(|\xi|)$.

The full L-P Decomposition is given as follows: given $u \in \mathcal{S}'$, $a = \psi_0(D)u + \sum_{j=1}^{\infty} \psi(2^{-j}D)u$. We will write $h = 2^{-j}$ as a shorthand.

Lemma 6.1

Suppose $\chi \in C_c^\infty(\mathbb{R}^n)$. Then for $u \in \mathcal{S}(\mathbb{R}^n)$, $\|\chi(hD)u\|_{L^\infty} \leq Ch^{-n/p} \|u\|_{L^p}$ and $\|\chi(hD)u\|_{L^p} \leq (2\pi)^{-n} \|\widehat{\chi}\|_1 \|u\|_p$.

Proof. Recall the following inequalities

- Holder's Inequality: $\|fg\|_1 \leq \|f\|_p \|g\|_q$ for $1/p + 1/q = 1$ for $1 \leq p \leq \infty$.
- Minkowski's Inequality: $\|f + g\|_p \leq \|f\|_p + \|g\|_p$ and

$$\left\| \int F(x, t) dt \right\|_p \leq \int \|F(\cdot, t)\|_p dt.$$

- Young's inequality: $\|f * g\|_p \leq \|f\|_1 \|g\|_p$.

We have

$$\begin{aligned} \chi(hD)u(x) &= \mathcal{F}^{-1}(\chi(h\xi)\widehat{u}(\xi)) = (2\pi)^{-n} \iint e^{i(x-y)\xi} \chi(h\xi) u(y) dy d\xi \\ &= (2\pi h)^{-n} \int \widehat{\chi}\left(\frac{x-y}{h}\right) u(y) dy \\ &\leq (2\pi h)^{-n} \frac{C}{h^n} \|\widehat{\chi}(\cdot/h)\|_q \|u\|_p \end{aligned}$$

Then,

$$\|\widehat{\chi}(\cdot/h)\|_q = \left(\int |\widehat{\chi}(y/h)|^q dy \right)^{1/q} = h^{n/q} \|\widehat{\chi}\|_q.$$

It follows that

$$|\chi(hD)u(x)| \leq Ch^{-n+n/q} \|u\|_p = Ch^{-n/p} \|u\|_p.$$

For the second inequality, note that $\chi(hD)u(x) = (2\pi h)^n \widehat{\chi}(\cdot/h) * u$. Applying Young's Inequality,

$$\|\chi(hD)u\|_p \leq \frac{1}{(2\pi h)^n} \|\widehat{\chi}(\cdot/h)\|_1 \|u\|_p \leq \frac{1}{(2\pi)^n} \|\widehat{\chi}\|_1 \|u\|_p.$$

□

Theorem 6.2

For $u \in L^p$, $1 \leq p \leq \infty$, $u \in C^{0,\gamma}(\mathbb{R}^n)$ if and only if for every $\chi \in C_c^\infty(\mathbb{R}^n \setminus 0)$, $\|\chi(hD)u\|_\infty \leq Ch^\gamma$.

Proof. We start with the forward direction. Note that

$$\begin{aligned} \chi(hD)u(x) &= \frac{1}{(2\pi h)^n} \int \widehat{\chi}((x-y)/h) u(y) dy \\ &= (2\pi)^{-n} \int \widehat{\chi}(y) u(x-yh) dy \\ &= (2\pi)^{-n} \int \widehat{\chi}(y) (u(x-yh) - u(x)) dy \end{aligned}$$

So it follows that

$$|\chi(hD)u(x)| \leq C \|u\|_{C^{0,\gamma}} \int |\widehat{\chi}(y)| (hy)^\gamma dy \leq C \|u\|_{C^{0,\gamma}} h^\gamma \int |\widehat{\chi}(y)| |y|^\gamma dy$$

□

and the last integral is bounded since $\widehat{\chi}$ is a Schwartz function, so it follows that $|\chi(hD)u(x)| \leq C \|u\|_{C^{0,\gamma}} h^\gamma$. □

7 February 9th, 2021

7.1 Fourier Transform proof of Morrey's Inequality

Recall the Littlewood-Paley decomposition: There exists $\psi_0 \in C_c^\infty(\mathbb{R}^n)$, $\psi \in C_c^\infty(\mathbb{R}^n \setminus \{0\})$ such that

$$1 = \psi_0(\xi) + \sum_{j=0}^{\infty} \psi(2^{-j}\xi).$$

From this, we have for $u \in \mathcal{S}'$,

$$u = \psi_0(D)u + \sum_{j=0}^{\infty} \psi(2^{-j}D)u.$$

More generally, for $a \in \mathcal{S}(\mathbb{R}^n)$, $a(D)u = \mathcal{F}^{-1}(a(\xi)\widehat{u}(\xi))$. We were proving the following theorem:

Theorem 7.1

For $u \in L^p$, $1 \leq p \leq \infty$, $u \in C^{0,\gamma}(\mathbb{R}^n)$ if and only if for every $\chi \in C_c^\infty(\mathbb{R}^n \setminus 0)$, $\|\chi(hD)u\|_\infty \leq Ch^\gamma$.

Proof. We proved the forward direction last time. We now show the converse.

Denote

$$\Lambda_\gamma(u) = \sup_{0 < h < 1} h^{-\gamma} (\|\psi(hD)u\|_\infty + \max_k \|\psi_k(hD)u\|_\infty)$$

where $\psi_k(\xi) = \xi_k \psi(\xi)$.

We have the hypothesis: $\|u\|_p + \Lambda_\gamma(u) < \infty$. We want to show that $\|u\|_{C^{\gamma,0}} \leq C(\|u\|_p + \Lambda_\gamma(u))$. We first bound $\|u\|_\infty$. Note that

$$\begin{aligned} \|u\|_\infty &\leq \|\psi_0(D)u\|_\infty + \sum_j \|\psi(2^{-j}D)u\|_\infty \\ &\leq \|\psi_0(D)u\|_\infty + \sum_j 2^{-j\gamma} \Lambda_\gamma(u) \\ &\leq C\|u\|_p + (2^\gamma - 1)^{-1} \Lambda_\gamma(u). \end{aligned}$$

Now, we bound the quotient term, $|u(x) - u(y)|/|x - y|^\gamma$. In other words, we want

$$|u(x) - u(y)| \leq C(\|u\|_p + \Lambda_\gamma(u))r^\gamma,$$

if $|x - y| \leq r$.

Note that

$$u(x) - u(y) = \psi_0(D)u(x) - \psi_0(D)u(y) + \sum_j (\psi(2^{-j}D)u(x) - \psi(2^{-j}D)u(y))$$

It is enough to prove that $|\psi_0(D)u(x) - \psi_0(D)u(y)| \leq Cr^\gamma\|u\|_p$ and

$$\sum_{j=0}^{\infty} |\psi(2^{-j}D)u(x) - \psi(2^{-j}D)u(y)| \leq Cr^\gamma \Lambda_\gamma(u).$$

Note that

$$\begin{aligned}
|\psi_0(D)u(x) - \psi_0(D)u(y)| &\leq \sup(\nabla(\psi_0(D)u))|x - y| \\
&\leq |x - y| \frac{1}{(2\pi)^n} \sup \int \nabla |\widehat{\psi}_0(x - y)| |u(y)| dy \\
&\leq |x - y| \frac{1}{(2\pi)^n} \|\nabla \widehat{\psi}_0\|_q \|u\|_p.
\end{aligned}$$

For the second inequality, we prove for both high frequency and low frequency estimates. For the high ones,

$$|\psi(hD)u(x) - \psi(hD)u(y)| \leq 2\|\psi(hD)u\|_\infty \leq 2h^\gamma \Lambda_\gamma(u).$$

For low frequencies,

$$\begin{aligned}
|\psi(hD)u(x) - \psi(hD)u(y)| &\leq Cr \max_k \|D_{x_k} \psi(hD)u\|_\infty \\
&= Crh^{-1} \max_k \|hD_{x_k} \psi(hD)u\|_\infty \\
&= Crh^{-1} \max_k \|\psi_k(hD)u\|_\infty \\
&\leq Crh^{-1} \max_k \|\psi_k(hD)u\|_\infty \\
&\leq Crh^{\gamma-1} \Lambda_\gamma(u).
\end{aligned}$$

Then, note that

$$\sum_{2^j \leq s} Cr2^{-j(\gamma-1)} \leq C'rs^{1-\gamma}$$

and

$$\sum_{2^j > s} C2^{-j\gamma} \leq C''s^{-\gamma}.$$

It follows that

$$\sum_{j=0}^{\infty} |\psi(2^{-j}D)u(x) - \psi(2^{-j}D)u(y)| \leq C\Lambda_\gamma(u)(rs^{1-\gamma} + s^{-\gamma}) \leq Cr^\gamma \Lambda_\gamma(u)$$

□

8 February 11th, 2021

8.1 Finishing Morrey's Inequality

The original statement of the theorem.

Theorem 13 (Morrey's Inequality)

Suppose $u \in L^p(\mathbb{R}^n)$, $\nabla u \in L^p(\mathbb{R}^n)$ and $n < p \leq \infty$. Then there exists $u^* \in C^{0,\gamma}(\mathbb{R}^n)$, with $\gamma = 1 - \frac{n}{p}$ such that $u = u^*$ almost everywhere and $\|u^*\|_{C^{0,\gamma}} \leq C(\|u\|_p + \|\nabla u\|_p)$.

Last time, we showed the following theorem:

Theorem 14

For $u \in L^p$, $1 \leq p \leq \infty$, $u \in C^{0,\gamma}(\mathbb{R}^n)$ if and only if for every $\chi \in C_c^\infty(\mathbb{R}^n \setminus \{0\})$, $\|\chi(hD)u\|_\infty \leq Ch^\gamma$.

Recall that

$$\Lambda_\gamma(u) = \sup_{0 < h < 1} h^{-\gamma} (\|\psi(hD)u\|_\infty + \max_k \|\psi_k(hD)u\|_\infty)$$

where $\psi_k(\xi) = \xi_k \psi(\xi)$. We proved this by showing that $\|u\|_{C^{0,\gamma}} \leq C(\|u\|_p + \Lambda_\gamma(u))$. We now show the complete proof of Morrey's Inequality.

Proof. It suffices to show that $\Lambda_\gamma(u) \leq C\|\nabla u\|_p$. Recall that for all $\chi \in C_c^\infty(\mathbb{R}^n)$, we showed that $\|\chi(hD)u\| \leq Ch^{-n/p}\|u\|_p$. Note that

$$\begin{aligned} \|\varphi(hD)hD_{x_j}u\|_\infty &\leq ch^{1-n/p}\|\nabla u\|_p \\ \Rightarrow \|\varphi_j(hD)u\|_\infty &\leq Ch^\gamma\|\nabla u\|_p. \end{aligned}$$

We would like to write $\psi \in C_c^\infty(\mathbb{R}^n \setminus \{0\})$, $\psi(\xi) = \sum \xi_j \chi_j(\xi)$ with $\chi_j \in C_c^\infty$. We can do this with $\sum \xi_j \frac{\xi_j}{|\xi|^2} \psi(\xi)$.

It follows that

$$\|\psi(hD)u\|_\infty \leq \sum_{j=1}^{\infty} \|\xi_j \chi_j(hD)u\|_\infty \leq Ch^\gamma\|\nabla u\|_p.$$

□

We can use this result to show regularity properties for solutions to PDEs. For example, one statement is as follows: suppose $u \in L^1$, $\Delta u = f \in C^{k,\gamma}$ for $0 < \gamma < 1$. We could show that $u \in C^{k+2,\gamma}$.

8.2 Final Comments about Sobolev Spaces

Definition 8.1. Suppose $U \subset \subset \mathbb{R}^n$ with $\partial U \in C^1$. Then $W_0^{1,p}(U) = \overline{C_c^\infty(U)}$ where the closure is respect to the $W^{1,p}$ norm.

Fact 8.2. $W_0^{1,p} = \{u \in W^{1,p}(U) : Tu = 0\}$, where $T : W^{1,p}(U) \rightarrow L^p(\partial U)$ linear and bounded and for $u \in W^{1,p}(U) \cap C(\bar{U})$, $Tu = u|_{\partial U}$.

Fact 8.3 (Poincare Inequality). Suppose $1 \leq p < n$ and $1 \leq q \leq p^* = \frac{np}{n-p}$. Then $\|u\|_q \leq C\|\nabla u\|_p$.

Theorem 15 (Poincare Inequality(v2))

For all $1 \leq p \leq \infty$, $\|u\|_p \leq C\|\nabla u\|_p$.

Proof. Suppose $p < n$. This follows from the version 1. Suppose $\infty > p \geq n$. In this case, take $q = n - \epsilon$. Then $q^* = \frac{(n-\epsilon)n}{\epsilon}$. Choose small enough ϵ so that $q^* \geq p$. Then, we apply Poincare 1: $\|u\|_p \leq \|u\|_{q^*} \leq C\|\nabla u\|_q \leq C\|\nabla u\|_p$. For $p = \infty$, the result follows from Morrey's inequality. \square

8.3 Duality

Recall the Riesz Representation Theorem for Hilbert Spaces:

Theorem 16 (Riesz Representation)

For $\Phi : H \rightarrow \mathbb{C}$ with a Hilbert space H , if $|\Phi(u)| \leq C\|u\|$, there exists $v \in H$ such that $\Phi(u) = \langle u, v \rangle$.

Fact 8.4. $H^{-s}(\mathbb{R}^n) = (H^s(\mathbb{R}^n))^*$: if $u \in H^{-s}(\mathbb{R}^n)$ and $v \in H^s(\mathbb{R}^n)$, $u \in H^s(\mathbb{R}^n)$, then $\langle u, v \rangle_{L^2} = \int u\bar{v}$ is well defined, and for any $\Phi : H^s(\mathbb{R}^n) \rightarrow \mathbb{C}$ such that $|\Phi(u)| \leq C\|u\|_{H^s}$, there exists $v \in H^{-s}$ such that $\Phi(u) = \langle u, v \rangle_{L^2}$.

Proof. First assume $u, v \in \mathcal{S}$. Then

$$\int u\bar{v} = (2\pi)^n \int \widehat{u}(\xi)\overline{\widehat{v}(\xi)} = (2\pi)^n \int \langle \xi \rangle^s \langle \xi \rangle^{-s} \widehat{v}(\xi)$$

so it follows that

$$|\langle u, v \rangle| \leq (2\pi)^n \|u\|_{H^s} \|v\|_{H^{-s}}.$$

Conversely, suppose we have Φ as above. Riesz implies that

$$\Phi(u) = \langle u, w \rangle_{H^s} = (2\pi)^{-n} \int \langle \xi \rangle^{2s} \widehat{u}\overline{\widehat{w}} = (2\pi)^{-n} \int \widehat{u}\overline{\langle \xi \rangle^{2s}\widehat{w}}$$

and we finish by setting $\widehat{v} = \langle \cdot \rangle^{2s}\widehat{w}$. \square

8.4 Duality on Bounded Domains

We define $H^{-1}(U) = \{u \in \mathcal{D}'(U) : \forall \varphi \in C_c^\infty(U), |u(\varphi)| \leq C\|\varphi\|_{H^1}\}$. The norm on H^{-1} is given by

$$\|u\|_{H^{-1}(U)} = \sup\{|u(\varphi)| : \varphi \in H_0^1, \|\varphi\|_{H^1} \leq 1\},$$

which is the usual operator norm, treating u as a linear functional on $H^1(U)$.

Example 8.5

We claim

$$H_0^1((0, \pi)) = \left\{ u(x) = \sum_{n=1}^{\infty} a_n \sin nx : \sum_{n=1}^{\infty} |a_n|^2 n^2 < \infty \sim \|u\|_{H^1} \right\}.$$

Then,

$$H^{-1}((0, \pi)) = \left\{ v(x) = \sum_{n=1}^{\infty} a_n \sin nx : \sum_{n=1}^{\infty} |a_n|^2 n^{-2} < \infty \right\},$$

where we take convergence in the sense of distributions.

Then,

$$\langle u, v \rangle = \sum a_n \bar{b}_n = \sum_{n=1}^{\infty} n a_n n^{-1} \bar{b}_n \leq \|u\|_{H^1} \|v\|_{H^{-1}}.$$

9 February 16th, 2021

9.1 Calculus of Variations: Minimizing Distance in the Plane

We start with a motivating example. Take points a, b in the x -axis, c, d in the y -axis. We wish to find a function $y = f(x)$ such that $f(a) = c$, $f(b) = d$ and the graph of f has the shortest length. Recall that

$$L(f) = \int_a^b (1 + f'(x)^2)^{1/2} dx.$$

We wish to minimize L over all paths from a to b . If f is a minimizer, then for all $\varphi \in C_c^\infty((a, b))$, $L(f + t\varphi)$ has a minimum at $t = 0$. This implies that $\frac{d}{dt}L(f + t\varphi)|_{t=0} = 0$ for all φ as above. Then,

$$\begin{aligned} \frac{d}{dt}L(f + t\varphi) &= \frac{d}{dt} \int_a^b (1 + (f + t\varphi)'^2)^{1/2} dx \\ &= \int_a^b \frac{\partial}{\partial t} [(1 + (f + t\varphi)'^2)^{1/2}] dx \\ &= \int_a^b \frac{\varphi'(x)(f'(x) + t\varphi'(x))}{(1 + (f'(x) + t\varphi'(x))^2)^{1/2}} dx \end{aligned}$$

Applying $t = 0$, we have

$$0 = \int_a^b \varphi'(x) \frac{f'(x)}{(1 + f'(x)^2)^{1/2}} dx$$

for all $\varphi \in C_c^\infty((a, b))$.

Integrating by parts, we get that

$$\int_a^b \varphi(x) \left(\frac{f'(x)}{(1 + f'(x)^2)^{1/2}} \right)' dx = 0.$$

This implies that

$$\frac{d}{dx} \left(\frac{f'(x)}{(1 + f'(x)^2)^{1/2}} \right) = 0,$$

with $f(a) = c$, $f(b) = d$.

We find that $f'(x) = \alpha$ so $f(x) = \alpha x + \beta$.

9.2 Calculus of Variations: Minimizing Area in \mathbb{R}^3

Take $U \subset \subset \mathbb{R}^2$, $\partial U \in C^1$. We wish to minimize the area of the graph with the condition that $f = g$ on ∂U .

We have

$$A(f) = \iint_U (1 + |\nabla f(x)|^2)^{1/2} dx.$$

We wish to minimize $A(f)$ over f satisfying $f = g$ on ∂U .

If f is a minimizer, $t \mapsto A(f + t\varphi)$, $\varphi \in C_c^\infty(U)$ has a minimum at $t = 0$. So,

$$\frac{d}{dt}A(f + t\varphi)|_{t=0} = 0.$$

Doing the same calculation as before, we have

$$\begin{aligned} \frac{d}{dt}A(f + t\varphi)|_{t=0} &= \int_U \frac{\partial}{\partial t}(1 + |\nabla f + t\nabla\varphi|^2)^{1/2} dx \\ &= \int_U \frac{\nabla\varphi \cdot \nabla f}{(1 + |\nabla f|^2)^{1/2}} dx \\ &= \int_U \varphi \left[\left(\frac{f_{x_1}}{(1 + |\nabla f|^2)^{1/2}} \right)_{x_1} + \left(\frac{f_{x_2}}{(1 + |\nabla f|^2)^{1/2}} \right)_{x_2} \right] dx. \end{aligned}$$

As before, this implies that

$$\left(\frac{f_{x_1}}{(1 + |\nabla f|^2)^{1/2}} \right)_{x_1} + \left(\frac{f_{x_2}}{(1 + |\nabla f|^2)^{1/2}} \right)_{x_2} = 0.$$

This is called the **Minimal Surface Equation**.

We will not solve this, but how could we do it? Consider $f \in H^1(U)$, and note that $Tf = g \in L^2(\partial U)$ is well-defined. If we take $m = \inf\{A(f) = f \in H^1(U), f|_{\partial U} = g\}$. Then, there exists $f_j \in H^1(U), f_j|_{\partial U} = g$ with $A(f_j) \rightarrow m$. Could we find $f_{j_k} \rightarrow f$?

9.3 Calculus of Variations: General Setup

Take $U \subset \subset \mathbb{R}^n$. Take $L : \mathbb{R}^n \times \mathbb{R} \times \bar{U} \rightarrow \mathbb{R}$ in C^∞ , written as $L(p, z, x)$. We introduce the functional $I[w] = \int_U L(Dw(x), w(x), x) dx$, with $w|_{\partial U} = g$.

Example 9.1

In the minimal surface problem, $L(p, z, x) = (1 + |p|^2)^{1/2}$.

We first derive an equation satisfied by the minimizer. As before, we have $I[w] = \int_U L(Dw, w, x) dx$, a minimizer. This implies that $\frac{d}{dt}I[w + t\varphi]|_{t=0} = 0$ for all $\varphi \in C_c^\infty(U)$. Then,

$$\begin{aligned} \int_U \frac{d}{dt} [L(Dw + tD\varphi, w + t\varphi, x)] dx|_{t=0} &= \int_U (D\varphi \cdot D_p L(Dw, w, x) + \varphi D_z L) dx \\ &= \int_U \left(- \sum (L_{p_j}(Dw, w, x))_{x_j} + D_z L(Dw, w, x) \right) \varphi dx \\ &\implies \boxed{- \sum_{j=1}^n (L_{p_j}(Dw, w, x))_{x_j} + D_z L(Dw, w, x) = 0}, \end{aligned}$$

the **Euler-Lagrange Equation**.

Example 9.2

Take $L(p, z, x) = |p|^2/2 - f(x)z$.

$$I[w] = \int_U (|\nabla w(x)|^2/2 - f(x)w(x)) dx.$$

Since $L_{p_j} = p_j$, the Euler-Lagrange equation is given by

$$- \sum (w_{x_j})_{x_j} - f(x) = 0 \implies -\Delta w = f, w|_{\partial U} = g.$$

We can generalize this as follows: If we take $L = |p|^2/2 + F(z)$ and $f(z) = F'(z)$. The Euler-Lagrange equation is then $-\nabla w = f(w)$. For example, if we take $f(z) = z^p$, $F(z) = \frac{z^{p+1}}{p+1}$.

We could also take non-constant coefficients: $L(p, z, x) = \frac{1}{2} \sum a_{ij}(x)p_i p_j - f(x)z$, where $a_{ij} = a_{ji}$.

Then, $L_{p_j} = \frac{1}{2} \sum_{i=1}^n a_{ij}(x)p_i$. Then, the Euler-Lagrange equation is given by

$$-\sum_{i,j=1}^n \partial_{x_j}(a_{ij} \partial_{x_i} w(x)) = f(x).$$

When $\sum a_{ij}(x)\xi_i \xi_j \geq c|\xi|^2$ for all $\xi \in \mathbb{R}^n$, $x \in \bar{U}$, this is solvable.

9.4 Existence of Minimizers

- Coercivity: There exists $\alpha, \beta \geq 0$ with $L(p, z, u) \geq \alpha|p|^q - \beta$, for $1 < q < \infty$, for all $z \in \mathbb{R}$, $x \in \bar{U}$.

The condition gives the following bound: $I[w] \geq \alpha \|Dw\|_q^q - \beta \mu(U)$. We can always set $\beta = 0$ by translating L by a constant. Taking $\mathcal{A} = \{w \in W^{1,q}(U) : u|_{\partial U} = g\}$, we minimize $I[w]$ over \mathcal{A} .

- Lower semicontinuity: Suppose we have $u_k \rightharpoonup u$ weakly in $W^{1,q}$. Then,

$$I[u] \leq \liminf I[u_k].$$

10 February 18th, 2021

Recall, we have $L : \mathbb{R}^n \times \mathbb{R} \times \bar{U} \rightarrow \mathbb{R}$ for $U \in \mathbb{R}^n$, $\partial U \in C^1$. We denote $L = L(p, z, x)$, $D_p L = (\partial_{p_1} L, \dots, \partial_{p_n} L)$, etc. We also defined

$$I[w] = \int_U L(Dw(x), w(x), x) dx, \quad w|_{\partial U} = g.$$

As an example, $L(p, x) = \frac{1}{2} \sum a_{ij}(x) p_i p_j - f(x)z$. Last time, we used the principle that if w is a minimizer, for every $\varphi \in C_c^\infty(U)$, $t \mapsto I[w + t\varphi]$ has a local minimum at $t = 0$. This implies that if w is a minimizer, L satisfies the Euler-Lagrange equation:

$$-\sum_{j=1}^n (L_{p_j}(Dw, w, x))_{x_j} + D_z L(Dw, w, x) = 0.$$

10.1 Second-Derivative Test

If $i'(0) = 0$, $i''(0) > 0$, then we have a local minimum at 0.

By definition $i(t) = I[w + t\varphi]$, where $\varphi \in C_c^\infty$. Recall that

$$i'(t) = \int \left(\sum \varphi_{x_j} \partial_{p_j} L(Dw + t\varphi, w + t\varphi, x) + \varphi \partial_z L(Dw + t\varphi, w + t\varphi, x) \right) dx.$$

Then,

$$i''(0) = \int \left(\sum_{i,j} \varphi_{x_j} \varphi_{x_i} \partial_{p_i p_j} L + \sum_j \varphi \varphi_{x_j} \partial_z \partial_{p_j} L + \varphi^2 L_{zz} \right) dx$$

If this is at least 0 for all φ , what do we get about L ? This makes sense for φ that is Lipschitz and 0 at the boundary. If we choose $\varphi(x) = \epsilon \rho(\frac{x\xi}{\epsilon}) \zeta(x)$, where $\zeta \in C_c^\infty(U)$ and ρ consists of triangles with slope ± 1 starting at 0. Then $|\rho'(x)| = 1$ almost everywhere.

Using this φ , we get $\varphi_{x_j} = \epsilon \rho(x\xi/\epsilon) \zeta'(x) + \xi_j \rho'(x\xi/\epsilon) \zeta(x) = \xi_j \rho'(x\xi/\epsilon) \zeta(x) + O(\epsilon)$ and

$$\begin{aligned} 0 &\leq \int_U \sum_{i,j} (\xi_i \xi_j \partial_{p_i p_j}^2 L) ((\rho')^2 \zeta^2) + O(\epsilon) \\ &\xrightarrow{\epsilon \rightarrow 0} \int_U \sum_{i,j} (\xi_i \xi_j \partial_{p_i p_j}^2 L) (\zeta^2) \end{aligned}$$

for any $\zeta \in C_c^\infty(U)$, so it follows that for all $\xi \in \mathbb{R}^n$, $\sum \xi_i \xi_j L_{p_i p_j}(Dw(x), w(x), x) \geq 0$. Hence, it is useful to assume convexity:

$$\sum_{i,j} \xi_i \xi_j L_{p_i p_j} L(p, z, x) \geq 0$$

for all $\xi \in \mathbb{R}^n$, $(p, z, x) \in \mathbb{R}^n \times \mathbb{R} \times U$. (

$$L(p + t\xi) = L(p) + t \sum \xi_j L_{p_j} L(p) + t^2 \int_0^1 (1-s) \sum \xi_i \xi_j L_{p_i p_j}(p + st\xi) ds$$

10.2 Convexity

For smooth L , convexity is the statement

$$\sum L_{p_i p_j}(p, z, x) \xi_i \xi_j \geq c |\xi|^2 \geq 0.$$

for all $\xi \in \mathbb{R}^n$.

Example 10.1

for $L = 1/2 \sum a_{ij}(x) p_i p_j$, $a_{ij} = a_{ji}$, convexity is that

$$\sum a_{ij}(x) \xi_i \xi_j \geq c |\xi|^2$$

We call this an Elliptic operator.

Example 10.2

For the minimal surface equation, $L = (1 + |p|^2)^{1/2}$. Note that $L_{p_i} = \frac{p_i}{(1 + |p|^2)^{1/2}}$.

$$L_{p_i p_j} = \frac{\delta_{ij}}{(1 + |p|^2)^{1/2}} - \frac{p_i p_j}{(1 + |p|^2)^{3/2}} = \frac{\delta_{ij}(1 + |p|^2) - p_i p_j}{(1 + |p|^2)^{3/2}}.$$

Then,

$$\begin{aligned} \sum L_{p_i p_j} \xi_i \xi_j &= \frac{1}{(1 + |p|^2)^{3/2}} \left(\sum |\xi|^2 (1 + |p|^2) - \sum \xi_i p_i \xi_j p_j \right) \\ &= (1 + |p|^2)^{3/2} (|\xi|^2 + |\xi|^2 |p|^2 - \langle \xi, p \rangle^2) \geq 0 \end{aligned}$$

This is not strictly convex, since as $p \rightarrow \infty$ our term goes to 0.

10.3 Existence of Minimizers

Recall our conditions:

- **Coercivity:** There exists $\alpha > 0, \beta \geq 0$ with $L(p, z, u) \geq \alpha |p|^q - \beta$, for some $1 < q < \infty$, for all $z \in \mathbb{R}, x \in \bar{U}$.

The condition gives the following bound: $I[w] \geq \alpha \|Dw\|_q^q - \beta \mu(U)$. We can always set $\beta = 0$ by translating L by a constant. Taking $\mathcal{A} = \{w \in W^{1,q}(U) : u|_{\partial U} = g\}$, we minimize $I[w]$ over \mathcal{A} .

- **Lower semicontinuity:** Suppose we have $u_k \rightharpoonup u$ weakly in $W^{1,q}$. Then,

$$I[u] \leq \liminf I[u_k].$$

As we will see, the coercivity leads to nice compactness results via Rellich-Kondrachov. How can we use lower semicontinuity? Assume \mathcal{A} is nonempty. Take $m = \inf_{w \in m \mathcal{A}} I[w]$. Then, we have $I[w_j] \rightarrow m$. Assuming coercivity, we have $\|Dw_j\|_q$ is bounded. If $w_0 \in \mathcal{A}$, then $\|w - w_0\|_q \leq \|Dw - Dw_0\|_q$ by the Poincare inequality. So it follows that $\|w_j\|_q \leq C$. From the Banach-Alaoglu Theorem, we have w_j is weakly compact in $W^{1,q}$. Passing to a subsequence, $w_j \rightharpoonup w$ in $W^{1,q}$. From lower semicontinuity, we have $I[w] \leq \liminf I[w_j] = m$. This implies that $I[w] = m$.

11 February 23rd, 2021

11.1 Weak Convergence

We have a Banach space B with dual B^* with $u : B \rightarrow \mathbb{C}$ linear in the dual if for all $x \in B$, $|u(x)| \leq C\|x\|_B$.

Theorem 17 (Banach - Alaoglu)

The unit ball $\{u \in B^* : \|u\|_{B^*} \leq 1\}$ is weak-* compact: if we have $\|u_j\|_{B^*} \leq 1$, then there exists a subsequence and $u \in B^*$ such that for every $x \in B$, $u_{j_k}(x) \rightarrow u(x)$.

Corollary 11.1

If B is reflexive, $(B^*)^* = B$, then $\{x : \|x\|_B \leq 1\}$ is weakly compact. Given $\|x_j\|_B \leq 1$, there exists $x \in B$ and a subsequence so that $u(x_{j_k}) \rightarrow u(x)$ for $u \in B^*$.

Example 11.2

Take $B = L^q(U)$ for $1 < q < \infty$. This is reflexive since $B^* = L^{q'}(U)$ with $q^{-1} + (q')^{-1} = 1$ for $1 < q' < \infty$.

Remark 11.3. For $u_j \in B$, we say $x_j \rightharpoonup x \in B$ iff for all $u \in B^*$, $u(x_j) \rightarrow u(x)$.

- If B reflexive and $x_j \rightharpoonup x$, then $\|x\|_B \leq \liminf \|x_j\|_B$. This is because $|x(u)| = \lim |x_j(u)| \leq \liminf \|u\|_{B^*} \|x_j\|_B$ and $\|x\|_B = \sup_{\|u\|_{B^*}=1} |x(u)|$.
- If B reflexive and $x_j \rightharpoonup x$, there exists C such that $\|x_j\|_B \leq C$. For every $u \in B^*$, $|x_j(u)| \leq C(u)$, which implies by the Uniform Boundedness Principle that $\|x_j\|_B \leq C$.
- (We don't assume B is reflexive) Suppose $V \subset B$ is a closed subspace. Then V is weakly closed. This is a special case of Mazur's Theorem.

Proof. We need to show that if $x_j \in V$, $x_j \rightharpoonup x \in B$, then $x \in V$. For $u \in B^*$, $u(x_j) \rightarrow u(x)$. So if $x \notin V$, we want to construct $u \in B^*$ so that $u(x_j) = 0$ and $u(x) = 1$.

Recall Hahn-Banach: If we have a subspace $\tilde{V} \subset B$ and $\tilde{\varphi} : \tilde{V} \rightarrow \mathbb{C}$ with $|\tilde{\varphi}(x)| \leq C\|x\|_B$, with $x \in \tilde{V}$, then there exists $\varphi \in B^*$ so that $\varphi|_{\tilde{V}} = \tilde{\varphi}$.

Take $\tilde{V} = V + \mathbb{C}x$. Define $\tilde{\varphi} : \tilde{V} \rightarrow \mathbb{C}$ and define $\tilde{\varphi}(y + \alpha x) = \alpha$, $y \in V$, $\alpha \in \mathbb{C}$. It suffices to check that it is bounded. We need $|\tilde{\varphi}(y + \alpha x)| \leq C\|y + \alpha x\|_B$. Suppose not - for every n , there exists y_n, α_n such that $|\alpha_n| = |\tilde{\varphi}(y_n + \alpha_n x)| > n\|y_n + \alpha_n x\|_B$.

Dividing by α_n , we get

$$1/n > \|y_n/\alpha_n + x\|_B.$$

But this would imply that $-y_n/\alpha_n \rightarrow x \notin V$, but V is closed. □

11.2 Calculus of Variations

We now move back to calculus of variations. We have $I[w] = \int_U L(Dw(x), w(x), x) dx$ with $U \in \mathbb{R}^n$, $\partial U \in C^1$. We have $L = L(p, z, x) \in C^\infty(\mathbb{R}^n \times \mathbb{R} \times \bar{U})$. We wish to minimize L under the constraint that $w|_{\partial U} = g$.

Recall that we introduced $i(t) = I[w + t\varphi]$, $\varphi \in C_c^\infty(U)$. If w is a minimizer of $I[w]$, then

- $i'(0) = 0$ for all φ implies that $-\sum_{j=1}^n \partial_{x_j}(\partial_{p_j} L(Dw, w, x)) + \partial_z L(Dw, w, x) = 0$ (Euler-Lagrange equation).
- If $i''(0) \geq 0$ for all φ , then we have the convexity condition: for all $\xi \in \mathbb{R}^n$, $\sum_{i,j} \partial_{p_i} \partial_{p_j} L(Dw, w, x) \xi_i \xi_j \geq 0$.

We introduced the conditions:

- Coercivity: there exists $1 < q < \infty$, $\alpha > 0$, $\beta \geq 0$ such that $L(p, z, x) \geq \alpha|p|^q - \beta$. This implies that $I[w] \geq \alpha \|Dw\|_q^q - \beta$.
- (Weak) Lower semicontinuity: If $u_j \rightharpoonup u$ and $Du_j \rightharpoonup u$ weakly in $L^q(U)$, then $I[u] \leq \liminf I[u_j]$.

Remark 11.4. Take $\mathcal{A} = \{u \in W^{1,q} : u|_{\partial U} = g\} \neq \emptyset$. If we put $m = \inf_{\mathcal{A}} I[w]$, there is a sequence $w_k \in \mathcal{A}$ such that $I[w_k] \rightarrow m$. Using weak compactness, we have a subsequence $w_{k_j} \rightharpoonup w$ in $W^{1,q}$ with $w \in \mathcal{A}$. Then $m \leq I[w] \leq \liminf I[w_{k_j}] = m$, so $I[w] = m$.

11.3 Getting around Lower Semicontinuity

Theorem 18

Suppose $L \geq -C$ and $p \mapsto L(p, z, x)$ is convex for all $(z, x) \in \mathbb{R} \times \bar{U}$. Then, for any $1 < q < \infty$, $w \mapsto I[w]$ is weakly lower semicontinuous in $W^{1,q}$: if $w_j \rightharpoonup w$ in L^q , $Dw_j \rightharpoonup Dw$ in L^q , then $I[w] \leq \liminf I[w_j]$.

Remark 11.5. Convexity implies that for all p_1, p_2 , $L(p_1) \geq L(p_2) + D_p L(p_2) \cdot (p_1 - p_2)$.

Proof. We assume that $u_k \rightharpoonup u$ in L^q and $Du_k \rightharpoonup Du$ in L^q . We have $\ell = \liminf I[u_k]$ and we want $I[u] \leq \ell$.

By taking a subsequence, we can say that $\ell = \lim I[u_k]$. By taking another subsequence, we can say $u_k \rightarrow u \in L^q$, since weak convergence implies that $\|u_k\|_q \leq C$ and $\|Du_k\|_q \leq C$ and using compactness of $W^{1,q}$ in L^q . By taking a subsequence we can use $u_k \rightarrow u$ almost everywhere [this is the Riesz-Fisher theorem]. By Egorov's Theorem, for every ϵ , there exists a set E_ϵ such that $m(U \setminus E_\epsilon) \leq \epsilon$ so that $u_k \rightarrow u$ uniformly on E_ϵ . Note that $m(U) < \infty$.

We define a set $F_\epsilon = \{x \in U : |u(x)| + |Du(x)| \leq \frac{1}{\epsilon}\}$. Then $m(U \setminus F_\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$. We define $G_\epsilon = E_\epsilon \cap F_\epsilon$, with $m(U \setminus G_\epsilon) \leq m(U \setminus E_\epsilon) + m(U \setminus F_\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$.

Without loss of generality, we can assume $L \geq 0$. Note that

$$\begin{aligned} I[u_k] &= \int_U L(Du_k, u_k, x) \\ &\geq \int_{G_\epsilon} L(Du_k, u_k, x) \\ &\geq \int_{G_\epsilon} L(Du, u_k, x) + D_p L(Du, u_k, x)(Du_k - Du) dx \end{aligned}$$

Then, $\lim \int_{G_\epsilon} L(Du, u_k, x) = \int_{G_\epsilon} L(Du, u, x)$ since $u_k \rightarrow u$ uniformly on G_ϵ and Du is uniformly bounded on G_ϵ . For the second term, $D_p L(Du, u_k, x) \rightarrow D_p L(Du, u, x)$ uniformly on G_ϵ . Then $Du_k \rightharpoonup Du$ in L^q . Then writing $\int (D_p L(Du, u_k, x) - D_p L(Du, u, x))(Du_k - Du) + D_p L(Du, u, x) \cdot (Du_k - Du)$, $Du_k - Du$ is bounded in L^q and $(D_p L(Du, u_k, x) - D_p L(Du, u, x))$ converges uniformly to 0, so the first term goes to 0. For the second term, $D_p L(Du, u, x)$ is bounded and $Du_k - Du$ converges weakly to 0.

It follows that

$$\ell = \liminf I[u_k] \geq \int_{G_\epsilon} L(Du, u, x) dx \xrightarrow{\epsilon \rightarrow 0} \int_U L(Du, u, x) dx = I[u].$$

Hence, $I[u] \leq \liminf I[u_k]$, as desired. □

12 February 25th, 2021

12.1 Existence of Minimizers

We proved last time that a convexity condition was sufficient for showing the weak lower semicontinuity condition.

Example 12.1

A simple example is $L(p, z, x) = \sum a_{ij}(x)p_i p_j$ where $a_{ij} = a_{ji}$ and $\sum a_{ij}(x)\xi_i \xi_j \geq \theta|\xi|^2$ for all $\xi \in \mathbb{R}^n$, $x \in \bar{U}$. In this case, $\mathcal{A} = \{u \in H^1(U) : u|_{\partial U} = g\}$, if $g \in H^{1/2}(\partial U)$. Then, we minimize $\int_U \sum a_{ij}(x)\partial_{x_j} u \partial_{x_i} u \, dx = 0$ with $u|_{\partial U} = g$.

Theorem 19 (Existence of Minimizers)

Suppose $p \mapsto L(p, z, x)$ is convex and $L(p, z, x) \geq \alpha|p|^q - \beta$, $\alpha > 0$, $\beta \geq 0$, $1 < q < \infty$. Suppose that $\mathcal{A} = \{w \in W^{1,q}(U) : w|_{\partial U} = g\} \neq \emptyset$ with $g \in L^q(\partial U)$, then there exists $u \in \mathcal{A}$ such that $I[u] = \min_{w \in \mathcal{A}} I[w]$.

Proof. We can assume without loss of generality that $\beta = 0$. Put $m = \inf_{\mathcal{A}} I[w] \neq \infty$. Choose a sequence $u_k \in \mathcal{A}$ such that $I[u_k] \rightarrow m$. Then $I[u_k] \geq \alpha \int |Du_k|^q$. This implies that $\|Du_k\|_{L^q} \leq C$.

Fix $w \in \mathcal{A}$ and note that $u_k - w \in W_0^{1,q}$. Recall the Poincaré Inequality: if $v \in W_0^{1,q}$, then $\|v\|_q \leq C\|Dv\|_q$. Hence

$$\|u_k\|_q \leq \|u_k - w\|_q + \|w\|_q \leq \|Du_k - Dw\|_q + \|w\|_q \leq \|Du_k\|_q + \|w\|_{W^{1,q}} \leq C'.$$

This implies that $\|u_k\|_{W^{1,q}} \leq C$. Hence, there exists a subsequence $u_k \rightharpoonup u$ in $W^{1,q}$. This means that $u_k - w \rightharpoonup u - w$ in $W^{1,q}$ but $u_k - w \in W_0^{1,q}$, a closed subspace in $W^{1,q}$ which implies that $u - w \in W_0^{1,q}$. Hence, $u \in \mathcal{A}$. So, it suffices to show that u is a minimizer.

From the convexity of $p \mapsto L(p, z, x)$, we have I is weakly lower-semicontinuous. In other words, $I[u] \leq \liminf_{k \rightarrow \infty} I[u_k] = \inf_{w \in \mathcal{A}} I[w]$. Hence, $I[u] = \inf_{w \in \mathcal{A}} I[w]$. \square

12.2 Uniqueness of Minimizers

Theorem 20 (Uniqueness of Minimizers)

Suppose $L = L(p, x)$ and there exists $\theta > 0$ such that for all $\xi \in \mathbb{R}^n$, $p \in \mathbb{R}^n$, $x \in \bar{U}$, we have $\sum L_{p_i p_j}(p, x)\xi_i \xi_j \geq \theta|\xi|^2$ (uniform convexity). Then, any minimizer of $I[u]$ is unique.

Proof. We have that $L(p, x) \geq L(q, x) \geq D_p L(q, x)(p - q) + \frac{\theta}{2}|p - q|^2$ from strict convexity (this follows from the Taylor remainder). Let u, \tilde{u} be minimizers of $I[w]$. Take $v = \frac{u + \tilde{u}}{2}$. From strict convexity, we have

$$I[u] \geq I[v] + \int D_p L(Du/2 + D\tilde{u}/2, x)(Du/2 - D\tilde{u}/2) + \theta/8|Du - D\tilde{u}|^2 \, dx.$$

Similarly,

$$I[\tilde{u}] \geq I[v] + \int D_p L(Du/2 + D\tilde{u}/2, x)(D\tilde{u}/2 - Du/2) + \theta/8 |Du - D\tilde{u}|^2 dx.$$

Then,

$$m = I[u]/2 + I[\tilde{u}]/2 \geq I[v] + \theta/8 \int |Du - D\tilde{u}|^2 dx \geq m + \theta/8 \int |Du - D\tilde{u}|^2 dx.$$

This would imply that $\theta/8 \int |Du - D\tilde{u}|^2 dx \leq 0$, which implies that $Du = D\tilde{u}$ almost everywhere but $u|_{\partial U} = \tilde{u}|_{\partial U}$ so $u = \tilde{u}$ almost everywhere. \square

12.3 The Euler-Lagrange Equation

Recall the example $L(p, x) = \sum a_{ij}(x)p_i p_j \geq \theta|p|^2$ with $a_{ij} = a_{ji}$. This has the Euler-Lagrange Equation:

$$\sum_{i,j=1}^n \partial_{x_i}(a_{ij}\partial_{x_j}u) = 0, \quad u|_{\partial U} = g.$$

For all $g \in H^{1/2}(\partial U)$ we can find $u \in H^1(U)$ such that the Euler-Lagrange equation weakly (in the sense of distributions). Furthermore, u is unique. We will show this next time.

Today, we show that we can solve the equation:

$$\sum_{i,j=1}^n \partial_{x_i}(a_{ij}\partial_{x_j}u) = f, \quad u|_{\partial U} = g, f \in H^{-1}(U)$$

with $u \in H^1(U)$. We can write

$$I[w] = \int \sum (a_{ij}(x)\partial_{x_i}w\partial_{x_j}w - f(x)w) dx$$

In this case $L(p, z, x) = \sum a_{ij}(x)p_i p_j - f(x)z$ for $z \in \mathbb{R}$, so this is not bounded below. However, $p \mapsto L(p, z, x)$ is convex. So it suffices to deal with the coercivity issue.

What we really need is that $I[w] \geq \int_U |Dw|^2 - \beta$.

Recall the Peter Paul inequality:

$$2ab \leq a^2/\epsilon + \epsilon b^2, \quad \forall \epsilon > 0.$$

Hence,

$$I[w] \geq \theta \int |Dw|^2 - \int |f||w| \geq \theta \int |Dw|^2 - \frac{1}{2\epsilon} \int |f|^2 - \frac{\epsilon}{2} \int |w|^2.$$

If we fix $w_0 \in H^1$ with $w_0|_{\partial U} = g \in H^{1/2}$, we have

$$I[w] \geq \theta \int |Dw|^2 - \frac{1}{2\epsilon} \int |f|^2 - \frac{\epsilon}{2} \left(\int |w - w_0|^2 + \int |w_0|^2 \right) \geq \theta \int |Dw|^2 - \frac{C}{2\epsilon} - \frac{\epsilon}{2} \int |Dw|^2.$$

It follows that

$$I[w] \geq \theta \int |Dw|^2 - C_\epsilon - \epsilon \int |Dw|^2 = (\theta - \epsilon) \int |Dw|^2 - C_\epsilon.$$

Choosing $\epsilon < \theta/2$, we have

$$I[w] \geq \frac{\theta}{2} \int |Dw|^2 - C_\epsilon \geq \alpha \int |Dw|^2 - C.$$

One problem: we assumed $f \in L^2$. How do we fix this? We solve 2 problems:

$$\sum (a_{ij}u_{x_j})_{x_i} = f \in H^{-1}, u|_{\partial U} = 0,$$

and $u \in H_0^1(U)$. Then, $\int fu \leq \|f\|_{H^{-1}}\|u\|_{H_0^1} \leq \frac{1}{\epsilon}\|f\|_{H^{-1}}^2 + \epsilon\|u\|_{H_0^1}^2$. For $u \in H_0^1$, $\|u\|_{H_0^1} \leq C\|Du\|_2$. Then, we apply the same argument.

If not, we take $v = u - \tilde{u}$ and we have

$$-\sum (a_{ij}v_{x_j})_{x_i} = 0, v \in H_0^1(U).$$

Otherwise, we multiply by v and we use the definition of weak solution. This implies that $\int \sum a_{ij}v_{x_i}v_{x_j}$, but it is also at least $\theta \int |Dv|^2$, which shows that it is exactly zero.

13 March 2nd, 2021

Today, we find conditions on the Lagrangian so that the the E-L condition holds:

$$-\sum(\partial_{p_j}L(Du(x), u(x), x))_{x_j} + \partial_zL(Du, u, x) = 0.$$

13.1 Euler-Lagrange Equation, continued

We will make the following assumptions: for all $p \in \mathbb{R}^n$, $z \in \mathbb{R}$, $x \in \bar{U}$,

- $|L(p, z, x)| \leq C(|p|^q + |z|^q + 1)$,
- $|D_pL|, |D_zL| \leq C(|p|^{q-1} + |z|^{q-1} + 1)$.

It is natural to consider the E-L equation in a weak sense, that is for all $v \in C_c^\infty(U)$,

$$\int \sum \partial_{p_j}Lv_{x_j} + \partial_zLv = 0.$$

The conditions imply that $|\partial_{p_j}L| \leq C(|Du|^{q-1} + |u|^{q-1}) \in L^{q'}(U)$, and the same with ∂_zL . This implies that our integral condition would make sense for $v \in W_0^{1,q}(U)$.

Definition 13.1. Suppose our assumptions from above hold(bounds on $|L|$ and $|DL|$) and $u \in \mathcal{A} = \{w \in W^{1,q}(U) : w|_{\partial U} = g\}$. We then say that the E-L equation holds weakly if for all $v \in W_0^{1,q}(U)$, we have

$$\int \sum \partial_{p_j}Lv_{x_j} + \partial_zLv = 0.$$

Theorem 21

Suppose $u \in \mathcal{A}$ is a minimizer for L satisfying the bounds. Then u is a weak solution to the Euler Lagrange equation.

Proof. Define $i(t) = I[u + tv]$ where $v \in W_0^{1,q}$. Let

$$\frac{i(t) - i(0)}{t} = \int_U \frac{L(Du + tDv, u + tv, x) - L(Du, u, x)}{t} dx$$

and call the integrand $L^t(x)$.

We have that $L^t(x) \xrightarrow{t \rightarrow 0} \sum L_{p_i}(Du, u, x)v_{x_i} + L_zv$ almost everywhere in x . We want to bound $|L^t(x)|$ by a function in L^1 so that we can apply the dominated convergence theorem.

Note that $f(\xi + t\eta) - f(\xi) = \eta \int_0^t f'(\xi + s\eta) ds$. This means that

$$L(Du + tDv, u + tv, x) - L(Du, u, x) = \int_0^t \sum L_{p_j}(Du + sDv, u + sv, x)v_{x_j} + L_zv ds.$$

Now, we bound this using our assumptions. Namely, recall that $|DL| \leq C(|p|^{q-1} + |z|^{q-1} + 1)$. It follows that

$$\begin{aligned} |L(Du + tDv, u + tv, x) - L(Du, u, x)| &\leq \int_0^t \sum |L_{p_j}(Du + sDv, u + sv, x)v_{x_j}| + |L_z v| ds \\ &\leq C \int_0^t (|Du + sDv|^{q-1} + |u + sv|^{q-1} + 1)(|Dv| + |v|) ds \\ &\leq C \int_0^t (1 + |Du|^{q-1} + |Dv|^{q-1} + |u|^{q-1} + |v|^{q-1})(|Dv| + |v|) ds \\ &\leq Ct (|Du|^{q-1}(|Dv| + |v|) + |u|^{q-1}(|Dv| + |v|) + |Dv|^q + |v|^q + 1) \end{aligned}$$

We would like to say

$$(|Du|^{q-1}(|Dv| + |v|) + |u|^{q-1}(|Dv| + |v|) + |Dv|^q + |v|^q + 1) \leq C (|Du|^q + |u|^q + |Dv|^q + |v|^q + 1).$$

We do this via Young's inequality: $ab \leq \frac{a^{q'}}{q'} + \frac{b^q}{q}$. This implies that

$$|Du|^{q-1}|Dv| \leq C(|Du|^{(q-1)q'} + |Dv|^q),$$

and doing this for the other product terms gives the desired inequality.

It follows that $|L^t(x)|$ is bounded by an L^1 function, so we can apply the dominated convergence theorem, which gives the result. \square

Remark 13.2. The converse is not necessarily true. However, we have the following theorem:

Theorem 22

Suppose $u \in \mathcal{A}$ is a weak solution to the Euler-Lagrange equation. If $(p, z) \mapsto L(p, z, x)$ is convex for all $x \in U$, then u is a minimizer.

Proof. From convexity, we have that $L(p, z, x) + D_p L(p, z, x) \cdot (q - p) + D_z L(p, z, x)(w - z) \leq L(q, w, z)$. Upon integrating, if we set $p = Du$, $q = Dw$, $z = u$, $w = w(x)$ for $w \in \mathcal{A}$:

$$I[u] + \int_U D_p(Du, u, x) \cdot (Dw - Du) + D_z L(Du, u, x) \cdot (w - u) dx \leq I[w].$$

But $w - u \in W_0^{1,q}$ and u weakly satisfies the equation so it follows that the integral is 0 and for every $w \in \mathcal{A}$, $I[u] \leq I[w]$. \square

14 March 4th, 2021

14.1 Regularity

We will make the assumption that $L(p, z, x) = L(p) - zf(x)$. We also assume that $|L(p)| \leq C(|p|^2 + 1)$ and $|D_p L(p)| \leq C(|p| + 1)$, $|D_p^2 L(p)| \leq C$. Finally, we assume strong convexity: $\sum L_{p_i p_j}(p) \xi_i \xi_j \geq \theta |\xi|^2$ for all $p, \xi \in \mathbb{R}^n$.

Example 14.1

$L(p) = \frac{1}{2}|p|^2$ is an example of a function which satisfies the above conditions.

Last time, we showed (with weaker assumptions) that if $I[u]$ is a minimum, then u satisfies the E-L equation weakly. With a convexity condition, we have the converse as well. We assume that $u|_{\partial U} = 0$ for simplicity.

Proposition 14.2

There exists a constant $C = C(L, n, U)$ so that $\|u\|_{H^1} \leq C\|f\|_2$.

Proof. We use the weak E-L with $v = u$. Namely,

$$\int_U \sum L_{p_j}(Du) u_{x_j} = \int_U f u.$$

Then, strict convexity implies that $(DL(p) - DL(0)) \cdot p \geq \theta |p|^2$, with $p = Du$. Hence,

$$\theta \int_U |Du|^2 \leq \int_U D_p L(Du) \cdot Du - \int_U DL(0) \cdot Du = \int_U f u,$$

where the second term is 0 by the divergence theorem. Then

$$\theta \int_U |Du|^2 \leq \int_U f u \leq \int_U \frac{f^2}{\epsilon} + \epsilon u^2 = \|f\|^2/\epsilon + \epsilon \|u\|^2.$$

Hence, $\|Du\|_2^2 \leq \frac{1}{\epsilon\theta} \|f\|^2 + \frac{\epsilon}{\theta} \|u\|^2$. By the Poincare inequality, $\|u\|_2 \leq C\|Du\|_2$, so it follows that by taking epsilon small enough. \square

14.2 Interior Regularity

Theorem 23

Suppose that $-\sum (L_{p_j}(Du))_{x_j} = f$ weakly, $f \in L^2$, $u \in H_0^1$ with the same bounds as before. Then $u \in H_{loc}^2(U)$.

Proof. Take $V \Subset W \Subset U$ open sets. Choose a function $\zeta \in C_c^\infty(W)$ and $\zeta = 1$ near V .

Choose $D_k^h u(x) = \frac{u(x+he_k) - u(x)}{h}$. If h is small enough then it is well defined on U . Note that for $v \in C_c^\infty(W)$, $\int u D_k^{-h} v = -\int v D_k^h u$. Then, we define $v = -D_k^{-h}(\zeta^2 D_k^h u) \in H_0^1(W) \subset H_0^1(U)$.

Then noting that $(D_k^{-h} v)_{x_i} = D_k^{-h}(v_{x_i})$,

$$\int_U \sum D_k^h(L_{p_i}(Du))(\zeta^2 D_k^h u)_{x_i} = -\int_U f D_k^{-h}(\zeta^2 D_k^h u).$$

$$\begin{aligned}
D_k^h(L_{p_i}(Du)) &= \frac{1}{h} \int_0^1 \sum_j L_{p_i p_j}(sDu(x + he_k) + (1-s)Du(x)) ds \cdot (D_{x_j}u(x + he_k) - D_{x_j}u(x)) \\
&= \sum_{j=1}^n a_{ij}^h(x) D_k^h u_{x_j}(x)
\end{aligned}$$

where

$$a_{ij}^h(x) = \int_0^1 L_{p_i p_j}(sDu(x + he_k) + (1-s)Du(x)).$$

So, we have

$$\int_U \sum_{j=1}^n a_{ij}^h(x) D_k^h u_{x_j} (\zeta^2 D_k^h u)_{x_i} = - \int_U f D_k^{-h} (\zeta^2 D_k^h u).$$

The left hand side is

$$\int \sum a_{ij}^h D_k^h u_{x_j} D_k^h u_{x_i} \zeta^2 + \sum a_{ij}^h D_k^h u_{x_j} 2\zeta \zeta_{x_i} D_k^h u.$$

The first term is bounded from below by $\theta |D_k^h Du|^2 \zeta^2$ by strict convexity. The second term is bounded by $-\int \zeta |D_k^h Du| |D_k^h u| \geq -\epsilon \int \zeta^2 |D_k^h Du|^2 - \frac{1}{\epsilon} |D_k^h u|^2$.

We end up with

$$\int \zeta^2 |D_k^h Du|^2 dx \leq C \int_W (f^2 + |D_k^h u|^2).$$

□

15 March 9th, 2021

15.1 Inner Regularity

Last time, we discussed regularity in the special case where $L = L(p) - f(x)z$. We also assumed that $|D_p^k L(p)| \leq C(1 + |p|^{2-k})$ and strict convexity: for all $p, \xi \in \mathbb{R}^n$, $\sum L_{p_i p_j}(p) \xi_i \xi_j \geq \theta |\xi|^2$.

We know that for $U \Subset \mathbb{R}^n$, $\partial U \in C^1$ and $f \in L^2$, for all $g \in H^{1/2}(\partial U)$, there exists u such that for all $v \in H_0^1(U)$,

$$\int_U \left(\sum L_{p_j}(Du) v_{x_j} - f(x)v \right) dx = 0, \quad u|_{\partial U} = g.$$

Theorem 15.1

Suppose that $-\sum (L_{p_j}(Du))_{x_j} = f$ weakly, $f \in L^2$, $u \in H_0^1$ with the same bounds as before. Then $u \in H_{loc}^2(U)$.

Remark 15.2. The main idea in the proof is that you can estimate derivatives with cutoff functions if you only need local results. This requires carefully choosing the width of the quotients to stay away from the boundary.

Proof. We had a function $D_k^h u(x) = \frac{u(x+he_k) - u(x)}{h}$, and we chose a ζ so that $\zeta \equiv 1$ in V and $\zeta \in C_c^\infty(W)$ where $V \subset W \subset U$.

Then, we defined $v = -D_k^{-h}(\zeta^2 D_k^h u)$. Last time, we showed that

$$\int_U \sum a_{ij}^h(x) D_k^h u_{x_j} (\zeta^2 D_x^h u)_{x_i} = \int_U f D_k^{-h}(\zeta^2 D_k^h u),$$

where we have the bounds

$$\theta |\xi|^2 \leq \sum a_{ij}^h(x) \xi_i \xi_j \leq C |\xi|^2.$$

Differentiating the expression, the LHS gives

$$\int \sum a_{ij}^h(x) (D_k^h u_{x_j} D_k^h u_{x_i}) \zeta^2 + \int \sum a_{ij}^h(x) D_k^h u_{x_j} 2\zeta \zeta_{x_i} D_k^h u.$$

The first term is bounded below by $\theta \int_U |D_k^h Du|^2 \zeta^2$. The second term is bounded below by $-C \int \zeta |D_k^h Du| |D_k^h u|$.

Now, the RHS is bounded above by

$$\int_W |f| |D_k^{-h} Du| \zeta^2 + \int_W |f| \zeta |D_k^h u|.$$

Note that

$$D_k^h u = \int_0^1 u_{x_k}(x + t h e_k) dt.$$

So it follows that $\int_W |D_k^h D_k^{-h} u| \zeta^2 \leq \int |D_k^h Du| \zeta^2$. It follows that

$$\begin{aligned} \int_W |f| |D_k^h Du| \zeta^2 + \int_W |f| \zeta |D_k^h u| &\leq \frac{2}{\epsilon} \int |f|^2 + \epsilon \int |D_k^h Du|^2 \zeta^2 + \epsilon \int \zeta^2 |D_k^h u|^2 \\ &\leq \frac{2}{\epsilon} \int |f|^2 + \epsilon \int \zeta^2 |D_k^h u|^2 + \epsilon \int |D_k^h Du|^2 \zeta^2. \end{aligned}$$

By using Peter Paul on the LHS bound and making $\epsilon \ll 1$, we obtain the inequality

$$\theta/2 \int |D_k^h Du|^2 \zeta^2 \leq 2/\epsilon \int |f|^2 + \epsilon \int \zeta^2 |D_k^h u|^2 \leq 2/\epsilon \int |f|^2 + \epsilon \int |Du|^2.$$

Finally, we claim that $\int |D_k^h w|^2 \leq C \Rightarrow w_{x_k} \in L_{loc}^2$. We know that $D_k^h w$ is bounded in L^2 for all h , so it is weakly compact. Hence, $D_k^{h_j} w \rightharpoonup v \in L^2$. It follows that

$$\int (D_k^h w) \varphi = - \int w D_k^{-h} \varphi \rightarrow - \int w \varphi_{x_k} = \int w_{x_k} \varphi,$$

which is the weak derivative. □

15.2 Higher Regularity

Take $f \equiv 0$ and $\int L_{p_i}(Du)v_{x_i} = 0$ for all $v \in H_0^1$. Take $w \in C_0^\infty(U)$ and set $v = -w_{x_k}$. We have

$$- \int \sum L_{p_i}(Du) \partial_{x_k}(w_{x_i}) dx.$$

But $u \in H_{loc}^2$, so it follows that

$$\int \sum_{i,j} L_{p_i p_j}(Du) u_{x_j x_k} w_{x_i} dx.$$

Setting $\tilde{u} = u_{x_k} \in H^1$. We get that $\sum \partial_{x_j}(a_{ij}(x) \partial_{x_i} \tilde{u}) = 0$ where $a_{ij} = L_{p_i p_j}(Du)$. But the bounded coefficients don't give a strong enough condition to prove the result.

Theorem 15.3

Suppose that $w \in H_{loc}^1(U)$, $\sum \partial_{x_j}(a_{ij}(x) \partial_{x_i} w) = 0$ weakly, and $\theta |\xi|^2 \leq \sum a_{ij}(x) \xi_i \xi_j \leq C |\xi|^2$. Then, there exists $\gamma > 0$ such that $w \in C_{loc}^{0,\gamma}$ (DeGiorgi-Nash, Moser). Applying with $w = \tilde{u} = u_{x_k}$ gives that $u \in C_{loc}^{1,\gamma}$, which implies that $a_{ij} = L_{p_i p_j}(Du) \in C^{0,\gamma}$. Finally, using Schauder estimates, we have that $u \in C^{2,\gamma}$.

16 March 11th, 2021

We follow the book *Grigis-Sjostrand: Microlocal Analysis for Differential Operators*.

16.1 Oscillatory Integrals

We denote $X \subset \mathbb{R}^n$ as an open set and $\mathcal{D}'(X) = \{u : C_c^\infty(X) \rightarrow \mathbb{C} : \forall K \subset\subset X, \exists C, N, \forall \varphi \in C_c^\infty(K), |u(\varphi)| \leq C \sup_{|\alpha| \leq N} |\partial^\alpha \varphi|\}$.

We wish to generalize expressions like $\delta_0(x) = \frac{1}{(2\pi)^n} \int e^{ix \cdot \xi} d\xi$. This is an "oscillatory integral" in the sense that we are integrating something that oscillates rapidly. This means that for all $\psi \in C_c^\infty$, $\delta_0(\psi) = \psi(0) = (2\pi)^{-n} \int \int e^{ix \cdot \xi} \psi(x) dx d\xi$.

We change the phase of $x \cdot \xi$ to a function $\varphi(x, \theta)$ where $x \in X$, $\theta \in \mathbb{R}^N$ so that $\varphi(x, \lambda\theta) = \lambda\varphi(x, \theta)$ for $\lambda > 0$.

The amplitude $(2\pi)^{-n}$ is generalized to $a(x, \theta)$, and we try to consider which properties we need so that we can define $I(a, \varphi) = \int_{\mathbb{R}^N} a(x, \theta) e^{i\varphi(x, \theta)} d\theta$ to be a distribution.

Example 16.1

Take $X = \mathbb{R}^n$, $N = n$. Define $P(\xi)$ to be a homogeneous polynomial satisfying $P(\xi) \neq 0$ whenever $\xi \neq 0$. For example, we could take $P(\xi) = |\xi|^2$. Define $\chi \in C_c^\infty(\mathbb{R}^n)$ so that $\chi \equiv 1$ near zero. Take $E(x) = (2\pi)^{-n} \chi(x) \int_{\mathbb{R}^n} \frac{1 - \chi(\xi)}{P(\xi)} e^{ix \cdot \xi} d\xi$. This doesn't converge, but the integrand is a smooth homogeneous function of degree $-m$ away from zero. If we define $D_x = \partial_x/i$, then

$$P(D)E(x) = \frac{1}{2\pi} \chi(x) \int_{\mathbb{R}^n} \frac{1 - \chi(\xi)}{P(\xi)} P(\xi) e^{ix \cdot \xi} + [P(D), \chi]u \int_{\mathbb{R}^n} \frac{1 - \chi}{R} e^{ix \cdot \xi} d\xi,$$

where $[P(D), \chi]$ is the commutator.

The first term is

$$\frac{1}{2\pi} \chi(x) \int e^{ix \cdot \xi} d\xi + (2\pi)^{-n} \chi(x) \int (-\chi(\xi)) e^{ix \cdot \xi} d\xi.$$

The second term is

$$\sum_{|\alpha| > 1} C \partial^\alpha X (2\pi)^{-n} \int (1 - \chi)/P \xi^\beta e^{ix \cdot \xi} d\xi.$$

The first term is $\delta_0(x)$, and the second is compactly supported. For the last term, for $x \neq 0$, $1/|x|^2 \langle x, \partial_\xi \rangle e^{ix \cdot \xi} = e^{ix \cdot \xi}$. The idea is that we can integrate by parts and the ∂_ξ derivatives will decay rapidly. Then, we can replace this term with something that decays rapidly so that we have $\delta_0(x) + K(x)$ for a compactly supported $K \in C_c^\infty(\mathbb{R}^n)$. This is called a "Parametrix" for $P(D)$.

In our above example, $\varphi(x, \theta) = x \cdot \theta$ and $\theta = \xi$, with $a(x, \xi) = \frac{1 - \chi}{P(\xi)}$. Our function satisfies the estimate

$$|\partial_\xi^\alpha ((1 - \chi)/P)| \leq C_\alpha \langle \xi \rangle^{-m - |\alpha|}.$$

16.2 General Theory: Amplitudes

Definition 16.2. $S_{\rho,\delta}^m(X \times \mathbb{R}^N) = \{a \in C^\infty(X \times \mathbb{R}^N) : \forall K \Subset X, \alpha \in \mathbb{N}^n, \beta \in \mathbb{N}^N, \exists C = C(K, \alpha, \beta) : |\partial_X^\alpha \partial_\xi^\beta a| \leq C \langle \xi \rangle^{m-\rho|\beta|+\delta|\alpha|}\}$.

In our previous example, when P is a homogeneous polynomial of degree q , $\frac{1-\chi(\xi)}{P(\xi)} \in S_{1,0}^{-q}$.

Remark 16.3. This only makes sense for $0 \leq \rho \leq 1$ and $0 \leq \delta \leq 1$. For $\rho > 1$. Then $|\partial^\alpha a| \leq C_{N,\alpha} \langle \xi \rangle^{-N}$. Suppose ξ has dimension $[in]$ and a has dimension $[in]^k$. $\partial_\xi^\alpha a$ has dimensions $[in]^{k-|\alpha|}$. On the other hand, $|\partial_\xi^\alpha a| \leq C \langle \xi \rangle^{m-\rho|\alpha|}$, which has dimension $[in]^{m-\rho|\alpha|}$. Then $k - |\alpha| = m - \rho|\alpha|$ so $k = m - (\rho - 1)|\alpha|$, but since α is arbitrary, the units of a are any negative number.

The same analysis would work for $\delta < 0$.

$S_{\rho,\delta}^m$ is a Frechet space, one that is generated by seminorms. Namely, note that $\|a\|_{K,\alpha,\beta} = \sup_{(x,\theta)} \langle \theta \rangle^{-m+\rho|\beta|+|\alpha|\delta} |\partial_x^\alpha \partial_\xi^\beta a|$. Then $a \in S_{\rho,\delta}^m$ if and only if for all $K \Subset X$, $\|a\|_{K,\alpha,\beta} < \infty$. The space is a Frechet space if it is complete with respect to this norm. This is also a metrizable space, with a metric defined in the obvious way.

Some properties:

- If $m \leq m'$, $\delta \leq \delta'$, $\rho \geq \rho'$, then $S_{\rho,\delta}^m \subset S_{\rho',\delta'}^{m'}$.
- We define $S^{-\infty}(X \times \mathbb{R}^N) = \{a \in C^\infty(X \times \mathbb{R}^N) : K \Subset X, \forall N, \exists C |\partial_x^\alpha \partial_\xi^\beta a| \leq C \langle \xi \rangle^{-N}\}$.
- $S^{-\infty}(X \times \mathbb{R}^N) = \bigcap_m S_{\rho,\delta}^m(X \times \mathbb{R}^N)$. We call this the residual space.

Example 16.4

Take $a \in C^\infty(X \times \mathbb{R}^N)$ and for $|\theta| \geq 1$, $\lambda > 0$, $a(x, \lambda\theta) = \lambda^m a(x, \theta)$. We claim that $a \in S_{1,0}^m$. If we differentiate, we have $\partial_\theta^\alpha a(x, \lambda\theta) = \lambda^{m-|\alpha|} \partial_\theta^\alpha a(x, \theta)$.

17 March 16th, 2021

17.1 Amplitudes of Oscillatory Integrals

We are making sense of $I(a, \varphi) = \int_{\mathbb{R}^N} a(x, \theta) e^{i\varphi(x, \theta)} d\theta$ on a distribution. We defined a class of functions for $m \in \mathbb{R}$, $\rho, \delta \in [0, 1]$. $S_{\rho, \delta}^m(X \times \mathbb{R}^N) = \{a \in C^\infty(X \times \mathbb{R}^N) : \forall K \Subset X, \alpha \in \mathbb{N}^n, \beta \in \mathbb{N}^N, \exists C = C(K, \alpha, \beta) : |\partial_X^\alpha \partial_\xi^\beta a| \leq C \langle \xi \rangle^{m - \rho|\beta| + \delta|\alpha|}\}$. These are called the **symbols** of order m and type (ρ, δ) . We write $S^m = S_{1, 0}^m$, which is the case where $|\langle \theta \rangle^{-m + |\beta|} \partial_x^\alpha \partial_\xi^\beta a| \leq C_{\alpha\beta}$.

Why are these called symbols? Suppose we have $P(x, 0)u = \sum_{|\alpha| \leq m} a_\alpha(x) D_k^\alpha u$. We can also write this as $(2\pi)^{-n} \int \sum_{|\alpha| \leq m} a_\alpha(x) \xi^\alpha e^{i(x-y)\xi} u(y) dy d\xi$, $u \in \mathcal{S}$. The current order of integration makes sense but we can also just consider the integral in $d\xi$, which is an oscillatory integral, whose integrand is $p(x, \xi)$ is the symbol of $P(x, D)$. Then $p \in S^m(X \times \mathbb{R}^n)$, which is of type $(1, 0)$.

17.2 A Cool Example

Suppose $f \in C^\infty(X \times \mathbb{R}^n; [0, \infty))$ homogeneous - $f(x, \lambda\theta) = \lambda f(x, \theta)$, $\lambda > 0$, $|\theta| \geq 1$. Define $a(x, \theta) = e^{-f(x, \theta)}$. Note that $0 \leq a \leq 1$. This is also a smooth function.

We claim that

$$\partial_x^\alpha \partial_\theta^\beta (e^{-f}) = \sum_{|\tilde{\alpha}| \leq |\alpha|, |\tilde{\beta}| \leq |\beta|} a_{\alpha\beta}(x, \theta) (\partial_x f)^{\tilde{\alpha}} (\partial_\theta f)^{\tilde{\beta}} e^{-f}.$$

We can estimate the bad term $(\partial_x f)^{\tilde{\alpha}} (\partial_\theta f)^{\tilde{\beta}} e^{-f}$. For this, we use Landau's inequality: If $g \in C^2(U)$, $g \geq 0$, for all $K \Subset U$, there exists C such that $|\nabla g(x)| \leq C \sqrt{g(x)}$, $x \in K$.

Proof. Note that $0 \leq g(x+y) = g(x) + \nabla g(x) \cdot y + O(|y|^2)$. This implies that $-\nabla g(x) \cdot y \leq g(x) + O(|y|^2)$. Taking $y = -\epsilon \nabla g(x)$, it follows that

$$\epsilon |\nabla g(x)| \leq g(x) + O(\epsilon^2 |\nabla g(x)|^2),$$

which implies the result. \square

We have that $f \geq 0$ so we have that $|\partial_x f| + |\partial_\theta f| \leq C f^{1/2}$, with $1 \leq \theta \leq 2$, $x \in K$. For $\lambda > 0$, note that

$$\lambda^{-1} \partial_x f(x, \lambda\theta) + \partial_\theta f(x, \theta) \leq C \lambda^{-1/2} f(x, \lambda\theta)^{1/2}.$$

It follows by taking $\tilde{\theta} = \lambda\theta$,

$$|\theta|^{-1/2} \partial_x f(x, \theta) + |\theta|^{1/2} \partial_\theta f(x, \theta) \leq C (f, \theta)^{1/2}, \quad |\theta| \leq 1, x \in K.$$

Now, we estimate

$$|(\partial_x f)^{\tilde{\alpha}} (\partial_\theta f)^{\tilde{\beta}} e^{-f}| \leq |\theta|^{|\tilde{\alpha}|/2} |\theta|^{-|\tilde{\beta}|/2} f^{1/2(|\tilde{\alpha} + \tilde{\beta}|)} e^{-f}.$$

Finally $f^k e^{-f} \leq k!$, which is some constant, so it follows that our term is bounded by $C_{\tilde{\alpha}\tilde{\beta}} |\theta|^{|\tilde{\alpha}|/2} |\theta|^{-|\tilde{\beta}|/2} \in S_{1/2, 1/2}^0(X \times \mathbb{R}^N)$.

17.3 Topology of the Symbol Space

Proposition 17.1

Suppose $\{a_j\}$ is bounded in $S_{\rho,\delta}^m$ and $a_j(x, \theta) \rightarrow a(x, \theta)$ for all $x, \theta \in X \times \mathbb{R}^N$. Then, $a \in S_{\rho,\delta}^m$ and $a_j \rightarrow a$ in $S_{\rho,\delta}^{m'}$ for all $m' > m$.

Proof. We first prove a lemma.

Lemma 17.2

Suppose $f \in C^2([- \epsilon, \epsilon])$. Then $|f'(0)| \leq 2\|f\|_{L^\infty}^{1/2}\|f''\|_{L^\infty}^{1/2} + (2/\epsilon + 1/2)\|f\|_{L^\infty}$, where $\|g\|_{L^\infty} = \sup_{|x| \leq \epsilon} |g|$.

Proof. $f(x) = f(0) + xf'(0) + x^2 + \int_0^1 (1-t)f''(tx) dt$. We have the estimate

$$|xf'(0)| \leq 2\|f\|_{L^\infty} + \frac{x^2}{2}\|f''\|_{L^\infty}.$$

Dividing by x , we we have $|f'(0)| \leq 2/x\|f\|_{L^\infty} + x/2\|f''\|_{L^\infty}$. Then, we take $x = \min(2\|f\|_{L^\infty}^{1/2}/\|f''\|_{L^\infty}^{1/2}, \epsilon) \leq 2\|f\|_{L^\infty}^{1/2}\|f''\|_{L^\infty}^{1/2} + (\frac{2}{\epsilon} + \frac{1}{2})\|f\|_{L^\infty}$, \square

From the lemma,

$$\|a'_j - a'_k\| \leq C\|a_j - a_k\|_{L^\infty}^{1/2}\|a''_j - a''_k\|_{L^\infty}^{1/2} + C\|a_j - a_k\|_{L^\infty}.$$

Hence, a Cauchy sequence in L^∞ implies Cauchy for higher symbols.

Example 17.3

Take $a = 1$, $a_j(\theta) = \chi(\theta/j)$. $a_j(\theta) \rightarrow a(\theta)$ for all θ . Do we have convergence of $a_j \rightarrow a$ in S^0 ? No! because $\|a_j - a\|_{L^\infty} = 1$. This is similar to the statement C_0^∞ is dense in L^p for $1 \leq p < \infty$.

Now, $\|\langle \theta \rangle^{-\delta}(a_j - a)\|_{L^\infty} \rightarrow 0$ as $j \rightarrow \infty$.

Define

$$b_j = \frac{\partial_x^\alpha \partial_\theta^\beta (a_j - a)}{\langle \theta \rangle^{m' - \rho|\beta| + \delta|\alpha|}} = \frac{1}{\langle \theta \rangle^{m' - m}} \frac{\partial_x^\alpha \partial_\theta^\beta (a_j - a)}{\langle \theta \rangle^{m - \rho|\beta| + \delta|\alpha|}}.$$

We know that $\partial_x^\alpha \partial_\theta^\beta (a_j - a)$ goes to 0 on compact sets, and $\langle \theta \rangle^{m-m'} \rightarrow 0$ as $|\theta| \rightarrow \infty$. Then, for all $\epsilon > 0$, there exists R_ϵ so that $|k_j| < \epsilon$ if $|\theta| > R_\epsilon$. On the other hand, for $|\theta| \leq R_\epsilon$, $x \in K$, $|k_j| < \epsilon$ if $j > J_\epsilon$. These two things imply that $|k_j| < \epsilon$ for $j > J_\epsilon$. \square

18 March 18th, 2021

18.1 Topology of the Symbol Space, continued

Recall that we wish to define a distribution in terms of the oscillatory integral $I(a, \varphi) = \int_{\mathbb{R}^N} u(x, \theta) e^{i\varphi(x, \theta)} d\theta$. We defined a class $a \in S_{\rho, \delta}^m(X \times \mathbb{R}^N)$ if for all $K \Subset X$, α, β , there exists C so that $|\partial_x^\alpha \partial_\theta^\beta a(x, \theta)| \leq C \langle \theta \rangle^{m - \rho|\alpha| + \delta|\beta|}$. This class is equipped with seminorms $\|a\|_{K, \alpha, \beta}$ in the obvious way and they generate the topology.

Last time, we proved the following: if $\{a_j\} \subset S_{\rho, \delta}^m$ is bounded ($\|a_j\|_{K, \alpha, \beta} \leq C_{K, \alpha, \beta}$), for all $(x, \theta) \in X \times \mathbb{R}^n$, $a_j(x, \theta) \rightarrow a(x, \theta)$. Then, $a \in S_{\rho, \delta}^m$ and $a_j \rightarrow a$ in $S_{\rho, \delta}^{m'}$ for $m' > m$.

Proposition 18.1

For every $m' > m$, $S^{-\infty}(X \times \mathbb{R}^N)$ is dense in $S_{\rho, \delta}^m$ in the $S_{\rho, \delta}^{m'}$ topology.

Proof. Write $\chi_j(\theta) = \chi(\theta/j)$ where χ is a bump function which is 1 on open set. Note that $\partial_\theta^\alpha \chi_j(\theta) = j^{-|\alpha|} \chi^{(\alpha)}(\theta/j)$, which is supported in $|\theta| \in [j, 2j]$, so we have $O(\langle \theta \rangle^{-|\alpha|})$. Hence, $\chi_j \in S_{1, 0}^0$ and moreover, $a_j := \chi_j a \in S_{\rho, \delta}^m$ uniformly: $\{a_j\}$ is a bounded sequence in $S_{\rho, \delta}^m$. On the other hand, $a_j \in S^{-\infty}$, so applying the result from last time implies the desired since $a_j \rightarrow a$ pointwise. \square

Remark 18.2. From Poincare, we say that $a \sim \sum_{j=0}^{\infty} a_j h^j$ (where the RHS doesn't converge) if for all N , there exists C so that $|a - \sum_{j=0}^{N-1} a_j h^j| \leq Ch^N$. This is like a Taylor's Theorem.

Theorem 18.3

Suppose $a_j \in S_{\rho, \delta}^{m_j}$, $m_j \rightarrow -\infty$, $m_0 \geq m_1 \geq \dots$. Then, there exists $a \in S_{\rho, \delta}^m$ such that for every k

$$a - \sum_{j=0}^{k-1} a_j \in S_{\rho, \delta}^{m_k}(X \times \mathbb{R}^N).$$

a is unique in the sense that if the equality holds for another \tilde{a} , then $a - \tilde{a} \in S^{-\infty}$.

We will write $a \sim \sum_{j=0}^{\infty} a_j$.

Proof. Suppose we have a sequence $\|\cdot\|_{k, \ell}$ is a sequence of seminorms defining the topology on $S_{\rho, \delta}^{m_k}$ (for k constant, $\|\cdot\|_{k, \ell}$ is a sequence of seminorms).

For every j , there exists $b_j \in S^{-\infty}$ so that $\|a_j - b_j\|_{\nu, \mu} \leq 2^{-j}$ for $0, \nu, \mu \leq j - 1$. This is okay because $S^{-\infty}$ is dense in $S_{\rho, \delta}^{m_j}$ in the topology of $S^{m_{nu}}$ for $\nu \leq j - 1$.

This implies that if we have $\sum_{j \geq k} a_j - b_j$, this converges in $S_{\rho, \delta}^{m_k}$ for all k . Now, we put $a = \sum_{j=0}^{\infty} a_j - b_j \in S_{\rho, \delta}^{m_0}$. Note that $a - \sum_{j < k} a_j = \sum_{j \geq k} a_j - \sum_{j=0}^{\infty} b_j = -\sum_{j < k} b_j + \sum_{j \geq k} a_j - b_j$. The first term is in $S^{-\infty}$ because it is a finite sum of things in $S^{-\infty}$. The second term is in $S_{\rho, \delta}^{m_0}$. \square

18.2 Phase Functions

We denote \mathbb{R}^N to be $\mathbb{R}^N \setminus \{0\}$.

Definition 18.4 (Non-degenerate Phase Function). A function $\varphi \in C^\infty(X \times \mathbb{R}^N)$ satisfying

- $\varphi(x, \lambda\theta) = \lambda\varphi(x, \theta)$, $\lambda > 0$.
- $Im(\varphi) \geq 0$.
- $d\varphi \neq 0$: $d\varphi = \sum \partial_{\theta_j} \varphi d\theta_j + \sum \partial_{x_j} \varphi dx_j$ or the gradient vector is different from zero for all x, θ .

Lemma 18.5

Suppose $m + k < -N$. Then, $a \mapsto I(a, \varphi)$ defines a continuous map between $S_{\rho, \delta}^m(X \times \mathbb{R}^N) \rightarrow C^k(X)$.

Proof. $I(a, \varphi) \in C(X)$ if $|a| \leq \langle \theta \rangle^{-N-\epsilon}$ i. e. $a \in S_{\rho, \delta}^m$ and $m < -N$. When we differentiate this k times, we get a function in $S_{\rho, \delta}^{m+k}$ which would map to $C(X)$ if $m + k < -N$. \square

Corollary 18.6

If $a \in S^{-\infty}$, this implies that $I(a, \varphi) \in C^\infty(X)$.

Theorem 18.7

Suppose $0 < \rho \leq 1$ and $0 \leq \delta < 1$. There exists a unique continuous map from $\bigcup_m S_{\rho, \delta}^m \ni a \mapsto I(a, \varphi) \in \mathcal{D}'(X)$, such that for $a \in S_{\rho, \delta}^m$, $m < -N$, $I(a, \varphi) = \int a(x, \theta) e^{i\varphi(x, \theta)} d\theta$. If $k \in \mathbb{N}$ and $m - k \min(\rho, 1 - \delta) < -N$, then the map $a \mapsto I(a, \varphi)$ is continuous as a map from $S_{\rho, \delta}^m \rightarrow \mathcal{D}'^{(k)}(X)$.

Remark 18.8. Recall that $u \in \mathcal{D}'^{(k)}(X)$ if for all $K \Subset X$, there exists c so that for all $\varphi \in C_c^\infty(K)$, $|u(\varphi)| \leq C \sup_{|\alpha| \leq k} |\partial^\alpha \varphi|$.

Proof. Uniqueness: this follows from the fact that $S^{-\infty}$ is dense in $S_{\rho, \delta}^m$ in $S_{\rho, \delta}^{m'}$ topology for all $m' > m$.

Existence: We sketch the proof and prove it next time.

- Use $d\varphi \neq 0$ to find a differential operator L so that $L^T e^{i\varphi} = e^{i\varphi}$.
- For $a \in S^{-\infty}$, from integration by parts, $I(a, \varphi)v = \int L^k(a, v) e^{i\varphi} d\theta dx$, which has stronger regularity.

\square

Example 18.9

Take $u(x) = \int_{\mathbb{R}^n} \langle \xi \rangle e^{ix \cdot \xi} d\xi$. Then $u(v) = \int_{\mathbb{R}^n \times \mathbb{R}^n} \langle \xi \rangle v(x) e^{ix \cdot \xi} dx d\xi$ where $v \in C_c^\infty$. Now, we construct L so that $L^T(e^{ix \cdot \xi}) = e^{ix \cdot \xi}$. This corresponds to $(\langle A, \partial_x \rangle + \langle B, \partial_\xi \rangle) e^{ix \cdot \xi} = e^{ix \cdot \xi}$. Then, we want $i\langle A, \xi \rangle + i\langle B, x \rangle = 1$.

19 March 30th, 2021

19.1 Oscillatory Integrals as Distributions

We had formal expressions $\int_{\mathbb{R}^N} a(x, \theta) e^{i\varphi(x, \theta)} d\theta$.

- $a \in S_{\rho, \delta}^M(X \times \mathbb{R}^N)$ if and only if $\partial_x^\alpha \partial_\xi^\beta a = O(\langle \xi \rangle^{m - |\beta|\rho + \delta|\alpha|})$.
- For a nondegenerate phase function, we have the conditions, φ is homogeneous of degree 1 in θ , $\text{Im}(\varphi) \geq 0$, and $d\varphi \neq 0$.

Lemma 19.1

If $a \in S_{\rho, \delta}^m$, $m + k < -N$, then $a \mapsto I(a, \varphi) = \int a e^{i\varphi}$ is continuous as a map from $S_{\rho, \delta}^m \mapsto C^k(X)$.

Theorem 19.2

Suppose $0 < \rho \leq 1$ and $0 \leq \delta < 1$. There exists a unique continuous map from $\bigcup_m S_{\rho, \delta}^m \ni a \mapsto I(a, \varphi) \in \mathcal{D}'(X)$, such that for $a \in S_{\rho, \delta}^m$, $m < -N$, it coincides with $I(a, \varphi) = \int a(x, \theta) e^{i\varphi(x, \theta)} d\theta$. If $k \in \mathbb{N}$ and $m - k \min(\rho, 1 - \delta) < -N$, then the map $a \mapsto I(a, \varphi)$ is continuous as a map from $S_{\rho, \delta}^m \rightarrow \mathcal{D}'^{(k)}(X)$.

Proof. We showed uniqueness last time. Namely, for all $a \in S_{\rho, \delta}^m$ there exists $a_j \in S_{\rho, \delta}^{-\infty}$ such that $a_j \rightarrow a$ in $S_{\rho, \delta}^{m'}$ for $m' > m$ so $I(a_j, \varphi)$ are uniquely determined by $\int a_j e^{i\varphi} \in C^\infty(X)$, so continuity implies the uniqueness.

For existence, we use the following lemma.

Lemma 19.3

Suppose that φ is a non-degenerate phase function. Then, there exists $a_j \in S_{1,0}^0$, $b_\ell \in S_{1,0}^{-1}$, $c \in S_{1,0}^{-1}$ such that if we define $L = \sum_j a_j(x, \theta) \partial_{\theta_j} + \sum_\ell b_\ell(x, \theta) \partial_{x_\ell} + c(x, \theta)$ so that ${}^tL(e^{i\varphi}) = e^{i\varphi}$.

Example 19.4

tL denotes the transpose of the operator when considered as a distribution. Namely, $Lv(u) = v({}^tLu)$. For example, ${}^t\partial_{x_j} = -\partial_{x_j}$. It is also easier to understand when taking it for test functions through integration.

Proof. Take $\chi \in C_c^\infty(\mathbb{R}^N)$ so that $\chi \equiv 1$ near 0. Define $\Phi = \sum |\frac{\partial \varphi}{\partial x_j}|^2 + |\theta|^2 \sum |\frac{\partial \varphi}{\partial \theta_j}|^2$. We have that Φ is a smooth function away from 0. It is homogeneous of degree 2 in θ , which can be understood by considering each component. This expression is nonzero away from 0. We can define the transpose

$${}^tL = \frac{1 - \chi(\theta)}{i\Phi} \sum |\theta|^2 \overline{\frac{\partial \varphi}{\partial \theta_j}} \partial_{\theta_j} + \sum \overline{\frac{\partial \varphi}{\partial x_j}} \partial_{x_j} + \chi(\theta).$$

We claim that ${}^tL(e^{i\varphi}) = e^{i\varphi}$. Note that $\varphi \mapsto i\Phi$. The first term is in S^0 , the second is in S^{-1} and $\chi \in S^{-\infty} \subset S^{-1}$. \square

We start with $u \in C_c^\infty(X)$. First, take $a \in S^{-\infty}$. Then, $I(a, \varphi) \in C^\infty(X)$, so we have

$$\begin{aligned} \langle I(a, \varphi), u \rangle &= \int_X \int_{\mathbb{R}^N} a(x, \theta) u(x) e^{i\varphi(x, \theta)} d\theta dx \\ &= \int_X \int_{\mathbb{R}^N} a u ({}^t L)^k (e^{i\varphi}) d\theta dx \\ &= \int_X \int_{\mathbb{R}^N} L^k(a u) e^{i\varphi} d\theta dx \end{aligned}$$

Now, suppose that $a \in S_{\rho, \delta}^m$ for $\rho > 0$ and $\delta < 1$. Then, $L^k(a u) \in S_{\rho, \delta}^{m-k \min(\rho, 1-\delta)}$, which means that for $\rho < 1$, $\delta > 1$, we gain decay. This gives a continuous map $(a, u) \mapsto S_{\rho, \delta}^m \times C_c^\infty(X)$.

Then, $\sup_{K \times \mathbb{R}^N} |L^k(a u) \langle \theta \rangle^{-m+k \min(\rho, 1-\delta)}| \leq C \|a\|_{\ell, K} \sum_{|\alpha| \leq k} \sup |\partial^\alpha u|$. This implies that if $a \in S_{\rho, \delta}^m$ and k is such that $m - k \min(\rho, 1 - \delta) < -N$, we can define

$$\langle I_k(a, \varphi), u \rangle = \int e^{i\varphi} L^k(a u) d\theta dx \leq C_{a, k, K} \sum_{|\alpha| \leq k} \sup |\partial^\alpha u|.$$

To define $I(a, \varphi)$ for $a \in S^m$ we need to show that if $m - k' \min(\rho, 1 - \delta) < -N$, then $I_k(a, \varphi) = I_{k'}(a, \varphi)$. We can prove this similarly to the uniqueness argument, since we have a sequence $a_j \rightarrow a$ in $S^{m'}$ for $m' > m$ and $I_k(a_j, \varphi) = I_{k'}(a_j, \varphi) = I(a, \varphi)$. \square

19.2 Some Remarks

- $I(a, \varphi) = \lim_{\epsilon \rightarrow 0} \int e^{i\varphi(x, \theta)} a(x, \theta) \chi(\epsilon \theta) d\theta$. This follows from the density of $a_\epsilon \rightarrow a$ in $S^{-\infty}$.
- Suppose $V = \sum a_j \partial_{\theta_j} + \sum b_\ell \partial_{x_\ell} + c$, $a_j \in S^0$, $b_\ell, c \in S^{-1}$. Then, $\iint e^{i\varphi} V(e^{i\varphi}) a u = \iint e^{i\varphi} V(a u)$, with the understanding that these are oscillatory integrals.
- We can use the above remark to prove things like smoothness of oscillatory integrals away from points. For example, take $(2\pi)^{-n} \int e^{ix\theta} d\theta = \delta_0(x)$. We can take $V = \frac{(1-\chi(x))x\partial_\theta}{|x|^2} \in S^0$ and it follows that $\int e^{ix\theta} d\theta \equiv 0$ away from 0.

20 April 1st, 2021

20.1 Oscillatory Integrals, continued

Recall that $\bigcup_m S_{\rho,\delta}^m \ni a \mapsto I(a, \varphi) \in \mathcal{D}'(X)$. We showed that for $a \in S^{-\infty}$, we had $I(a, \varphi) = \int a(x, \theta) e^{i\varphi(x, \theta)} d\theta \in C^\infty(X)$, with the assumption that φ is a non-degenerate phase function.

We mentioned last time that if we have $\iint {}^tL(e^{i\varphi})au d\theta dx = \iint e^{i\varphi} L(au) d\theta dx$ for any $u \in C_c^\infty(X)$ where $L = \sum a_j \partial_{\theta_j} + \sum b_k \partial_{x_k} + c$, $a_j \in S^0$, $b_k \in S^{-1}$, $c \in S^{-1}$.

Definition 20.1. Given a phase function φ , define $C_\varphi = \{(x, \theta) : d_\theta \varphi = 0\}$.

If we take $\delta_0(x)$, $\varphi = x \cdot \theta$, then $C_\varphi = \{(0, \theta) : \theta \in \mathbb{R}^n\}$.

Lemma 20.2

Suppose $a \in S_{\rho,\delta}^m$, $0 < \rho, \delta < 1$, $a \equiv 0$ in a conic neighborhood C_φ . Then $I(a, \varphi) \in C^\infty(X)$.

Definition 20.3. A conic neighborhood of $\theta_0 \in \mathbb{R}^N$ is a set θ satisfying $|\frac{\theta}{|\theta|} - \frac{\theta_0}{|\theta_0|}| < \epsilon$ for some ϵ .

Proof. Claim: there exists $L = \sum a_j \partial_{\theta_j} + c$, $a_j \in S^0$, $c \in S^{-1}$ such that ${}^tL(e^{i\varphi}) = (1-b)e^{i\varphi}$, $b \in S^0$ with $\text{supp } b \cap \text{supp } a = \emptyset$.

Namely, choose b , which is homogeneous of degree 0, $|\theta| > 1$ in a conic neighborhood of C_φ and $b = 0$ on the support of a . We can simply take $b(x, \theta)$, $\theta \in S^{N-1}$ so that $b = 1$ near $C_\varphi \cap S^{N-1}$. If $\text{supp } b$ is close enough to C_φ , this implies the disjoint supports by the assumption that $a \equiv 0$ on the conic neighborhood of C_φ .

Then, we construct ${}^tL = (1-b) \frac{1}{|\varphi_\theta|^2} \langle \varphi_\theta, \partial_\theta \rangle$. This is fine because $b \equiv 1$ on the set where φ_θ vanishes. We also know that $(1-b)\partial^\alpha a = \partial^\alpha a$ and $\partial^\alpha b \partial^\beta a = 0$. This implies that $I(a, \varphi) = I(L^k a, \varphi)$ for all k where we have $L(au) = (La)u$. Furthermore, note that $L^k a \in S^{m-k \min(\rho, 1-\delta)}$, so we can make this arbitrarily small for large k , which implies that $I(a, \varphi) \in C^m(X)$ for all m . □

Theorem 20.4

$\text{sing supp } I(a, \varphi) \subset \pi(C_\varphi)$ where $\pi : X \times \mathbb{R}^N \rightarrow X$, $\pi(x, \theta) = x$.

Remark 20.5. Recall that for $u \in \mathcal{D}'(X)$, $\text{sing supp } u = \{x : \exists U = \text{nbhd}(x), a|_U \in C^\infty(U)\}^c$.

Proof. Suppose $x_0 \notin \pi(C_\varphi)$. Then, there exists $\psi(x)$ such that $\psi(x_0) = 1$, $\text{supp } \psi \cap \pi(C_\varphi) = \emptyset$ (this exists because C_φ is closed, so its projection is closed). Then $\psi I(a, \varphi) = I(\psi a, \varphi)$. Now, $\text{supp } \psi a \cap C_\varphi = \emptyset$. Now, we apply the lemma. □

Some examples

- For $f \in C^\infty(X)$, $\text{Im}(f) \geq 0$ and $f(x) = 0$ implies that $df(x) \neq 0$ (this implies that $\{x; f(x) = 0\}$ is a C^∞ hypersurface). Define $u(x) = \int_0^\infty e^{if(x)\tau} d\tau$. Strictly

speaking, this should be $\int_0^\infty \chi(\tau)e^{if(x)\tau} d\tau + \int_{\mathbb{R}}(1 - \chi(\tau))e^{if(x)\tau} d\tau$, which is now $I(1 - \chi, f(x)\tau) \in S^0$.

Note that $\int_0^\infty e^{i(f(x)+i\epsilon)\tau} d\tau = \frac{i}{f(x)+i\epsilon}$ converges to $u(x) \in \mathcal{D}'(X)$. It follows that $u(x) = i(f(x))^{-1}$ and $C_\varphi = \{(x, \tau) : f(x) = 0\}$.

- We wish to solve $(\partial_t^2 - \Delta)u = 0$, $u|_{t=0} = f$, $\partial_t u|_{t=0} = 0$, $f \in S'(\mathbb{R}^n)$. Then, $u(t, x) = (2\pi)^{-n} 1/2 \iint \sum_{\pm} e^{i(x-y)\xi \pm |\xi|t} f(y) dy d\xi$. We can check this is a solution to the wave equation.

Define $U(t, x, y) = (2\pi)^{-n} \int \frac{1}{2} \sum_{\pm} e^{i(x-y)\xi \pm |\xi|t} d\xi \in \mathcal{D}(\mathbb{R} \times \mathbb{R}^n)$, since we can check directly that $\varphi_{\pm}(t, x, y, \xi)$ is a phase function. Note that $C_{\varphi_{\pm}} = \{|x - y| = t\}$, which gives the light cone.

20.2 Fourier Integral Operators and Pseudodifferential Operators

Suppose that $\varphi \in C^\infty(X \times Y \times \mathbb{R}^N)$, $X \subset \mathbb{R}^n$, $Y \subset \mathbb{R}^m$ is a phase function in all variables. Namely, $X \times Y$ is the "old X " and $a \in S_{\rho, \delta}^m(X \times Y \times \mathbb{R}^N)$, $\rho > 0$, $\delta < 1$. We obtain $K(x, y) = \int a(x, y, \theta)e^{i\varphi(x, y, \theta)} d\theta \in \mathcal{D}'(X \times Y)$. K defines $A : C_c^\infty(Y) \rightarrow \mathcal{D}'(X)$ via $u \in C_c^\infty(Y)$, $c \in C_c^\infty(X)$, and $\langle Au, v \rangle := \langle K, v \otimes u \rangle$, where $v \otimes u(x, y) = v(x)u(y) \in C_c^\infty(X \times Y)$.

Formally, we write $Au(x) = \iint a(x, y, \theta)e^{i\varphi(x, y, \theta)}u(y) dy d\theta$ which is understood as an oscillatory integral when paired with v . This is called the **Fourier Integral Operator**.

For $\theta \in \mathbb{R}^n$, $X = Y \subset \mathbb{R}^n$, $\varphi(x, y, \theta) = (x - y) \cdot \theta$, we have a **Pseudodifferential Operator**.

Suppose we have a differential operator $P(x, D) = \sum_{|\alpha| \leq m} a_\alpha(x)D_x^\alpha$, $a - \alpha \in C^\infty(X)$. We can write

$$Pu(x) = (2\pi)^{-n} \iint \sum a_\alpha(x)\xi^\alpha u(y) dy d\xi.$$

Then, $a(x, \xi) = \sum a_\alpha \xi^\alpha \in S_{1,0}^m(X \times \mathbb{R}^n)$.

Theorem 20.6

Suppose A is an FIO, we have the following:

- If for every $x \in X$, $(y, \theta) \mapsto \varphi(x, y, \theta)$ is a phase function, then $A : C_c^\infty(Y) \rightarrow C^\infty(X)$.
- If for every $y \in Y$, $(x, \theta) \mapsto \varphi(x, y, \theta)$ is a phase function, then $a : \mathcal{E}'(Y) \rightarrow \mathcal{D}'(X)$.

21 April 6th, 2021

21.1 Operators on Oscillatory Integrals

Take $X \subset \mathbb{R}^n$, $Y \subset \mathbb{R}^m$, $a \in S_{\rho,\delta}^m(X \times Y \times \mathbb{R}^N)$, $\varphi = \varphi(x, y, \theta)$ a non-degenerate phase function. Assuming $\rho > 0$, $\delta < 1$, this gives a distribution $K = I(a, \varphi) \in \mathcal{D}'(X \times Y)$. Furthermore, this defines a continuous operator $A : C_c^\infty(Y) \rightarrow \mathcal{D}'(X)$ given for $u \in C_c^\infty(Y)$, $v \in C_c^\infty(X)$ by

$$\langle Au, v \rangle = \langle K, u \otimes v \rangle$$

where $u \otimes v = u(x)v(y) \in C_c^\infty(X \times Y)$. These are called **Fourier Integral Operators**.

- $K_\pm(t, x, y) = (2\pi)^{-n} \int e^{i\langle \xi, x-y \rangle} e^{\pm i|\xi|t}$ for $t \in \mathbb{R}$, $x, y \in \mathbb{R}^n$.
- For $X = Y$, $\varphi(x, y, \xi) = \langle x - y, \xi \rangle$, this gives the **Pseudo-differential Operators**.

Remark 21.1. The map $a \mapsto I(a, \varphi) \in \mathcal{D}'(X \times Y)$ is continuous, so that for $u, v \in C_c^\infty(W)$, $W \Subset X \times Y$,

$$|\langle K, u \otimes v \rangle| \leq C \sum_{|\alpha| \leq N, |\beta| \leq N} \sup |\partial_x^\alpha \partial_y^\beta (uv)|.$$

Theorem 21.2

Suppose A is an FIO, we have the following:

- If for every $x \in X$, $(y, \theta) \mapsto \varphi(x, y, \theta)$ is a non-degenerate phase function ($d_{y,\theta}\varphi \neq 0$), then $A : C_c^\infty(Y) \rightarrow C^\infty(X)$.
- If for every $y \in Y$, $(x, \theta) \mapsto \varphi(x, y, \theta)$ is a non-degenerate phase function ($d_{x,\theta}\varphi \neq 0$), then $a : \mathcal{E}'(Y) \rightarrow \mathcal{D}'(X)$.

Example 21.3

For a pseudo-differential operator, we have $\varphi = \langle x - y, \xi \rangle$ and $(y, \xi) \mapsto \langle x - y, \xi \rangle$ and $(x, \xi) \mapsto \langle x - y, \xi \rangle$ are non-degenerate.

Proof. Take $\Phi = |d_y\varphi|^2 + |\theta|^2 |d_\theta\varphi|^2$, which is nonzero for $X \times Y \times \mathbb{R}^N$. This is homogeneous of degree 2. Let χ be the usual bump function and set

$${}^tL = \frac{1 - \chi}{i\Phi} (\langle \partial_y\varphi, \partial_y \rangle + |\theta|^2 \langle \partial_\theta\varphi, \partial_\theta \rangle) + \chi.$$

Then, $L = \langle A, \partial_y \rangle + \langle B, \partial_\theta \rangle + c$ for $A \in S^{-1}$, $B \in S^0$, $c \in S^{-\infty} \subset S^{-1}$. We showed that for operators of this form, we can integrate by parts. Namely,

$$\langle Au, v \rangle = \int e^{i\varphi} v(x) L^k(au) dx dy d\theta.$$

Then, $L^k \in S^{m-k \min(\rho, 1-\delta)}$. It follows that $Au = \int e^{i\varphi} L^k(au) dy d\theta \in C^\infty(X)$.

For the second part, note that for any $A : C_c^\infty(Y) \rightarrow \mathcal{D}'(X)$, we define ${}^tA : C_c^\infty(X) \rightarrow \mathcal{D}'(Y)$ by $\langle {}^tAv, u \rangle = \langle Au, v \rangle$. If A is defined using $K_A \in \mathcal{D}'(X \times Y)$, this implies that tA

is defined by $K_{t_A}(y, x) = K_A(x, y)$. This means that if $K_A = I(a, \varphi)$, then $K_{t_A} = I(\tilde{a}, \tilde{\varphi})$ where $\tilde{a}(y, x, \theta) = a(x, y, \theta)$ and $\tilde{\varphi}(y, x, \theta) = \varphi(x, y, \theta)$. But now, the condition that $(x, \theta) \mapsto \varphi(x, y, \theta)$ is non-degenerate implies that $K_{t_A} : C_c^\infty(X) \rightarrow C^\infty(Y)$. It follows that $A : \mathcal{E}'(Y) \rightarrow \mathcal{D}'(X)$. For $u \in \mathcal{E}'(Y)$, $v \in C_c^\infty(X)$, we define $\langle Au, v \rangle = \langle {}^tAv, u \rangle$. \square

21.2 Method of Stationary Phase

We defined $C_\varphi = \{(x, \theta) : \varphi_\theta = 0\}$. We proved that $\text{sing supp } I(a, \varphi) \subset \pi(C_\varphi)$ where $\pi(x, \theta) = x$. One could ask, when do we have equality? To do this, we need to "evaluate" our oscillatory integrals. This is the **Method of Stationary Phase**.

Define

$$I(\lambda) = \int e^{i\lambda\varphi(x)} a(x) dx,$$

for $a \in C_c^\infty(\mathbb{R})$, $\varphi \in C^\infty(\mathbb{R}, \mathbb{R})$. We could also take

$$J(\lambda) = \int e^{-\lambda\psi(x)} a(x) dx.$$

The study of the first is the stationary phase method and the study of the second is the steepest descent method.

21.3 Steepest Descent

Recall from lecture 1, we were stuck on the integral $\int_0^{2\pi} e^{-\frac{1}{2\epsilon}|\xi|^2 \cos^2 \theta} d\theta$ as $\epsilon \rightarrow 0$. The Method of Steepest Descent will give us the asymptotic expansion of this.

For $a \in C_c^\infty$, ψ has a unique non-degenerate minimum.

$\int e^{-\lambda\psi(x)} a(x) dx = e^{-\lambda\psi(x_0)} ((2\pi\lambda\psi''(x_0))^{-1/2} a(x_0) + b_1\lambda^{-1/2-1} + b_2\lambda^{-1/2-2} + \dots)$. This means that for all N , there exists C such that

$$\left| \int e^{-\lambda\psi(x)} a(x) dx - e^{-\lambda\psi_0(x_0)} (2\pi\lambda\psi''(x_0))^{-1/2} a(x_0) + b_1\lambda^{-1/2-1} + b_2\lambda^{-1/2-2} + \dots \right| \leq C e^{-\lambda\psi(x_0)} \lambda^{-N-5/2}.$$

If we take $\chi \in C_c^\infty(\mathbb{R})$, χ with support close to x_0 ,

$$\left| \int e^{-\lambda\psi(x)} (1 - \chi(x)) a(x) dx \right| \leq e^{-\lambda(\psi(x_0) + \epsilon)}.$$

By Taylor's Formula, $\psi(x) = 1/2(x - x_0)^2\psi_1(x)$, where $\psi(x_0) = \psi''(x_0)$. Then, take $y = y(x) = (x - x_0)(\psi_1(x))^{1/2}$ for x near x_0 . Since $y'(x_0) \neq 0$, we can write $x = x(y)$, for y near 0. Finally, $\varphi(x(y)) = y^2/2$, so we have

$$\int e^{-\lambda\psi(x)} \chi(x) a(x) dx = \int e^{-1/2\lambda y^2} b(y) dy$$

where $b(y) = \chi(x(y)) a(x(y)) |dx/dy|$. We know that $b(0) = a(x_0)/(\psi''(x_0))^{1/2}$. We are reduced to studying

$$\int e^{-1/2\lambda y^2} b(y) dy.$$

Then, $\int u\bar{v} = (2\pi)^{-1} \int \widehat{u\bar{v}}$, Plancherel's formula. Then, $u = b(y)$, $v = e^{-1/2\lambda y^2}$. The Fourier transform of $e^{-1/2\lambda y^2} = \frac{\sqrt{2\pi}}{\lambda^{1/2}} e^{-\xi^2/2\lambda}$, so it follows that

$$\int e^{-1/2\lambda y^2} b(y) dy = (2\pi\lambda)^{-1/2} \int e^{-\xi^2/2\lambda} \widehat{b}(\xi) d\xi.$$

We wish to consider this as $\lambda \rightarrow \infty$. By expanding the Taylor series of the exponential term, we have

$$(2\pi\lambda)^{-1/2} \sum \int \frac{1}{k!} (-1/2\lambda\xi^2)^k \widehat{\beta}(\xi) d\xi = (2\pi\lambda)^{-1} \sum \frac{(-1/2\lambda)^k}{k!} \int \xi^{2k} \widehat{b}(\xi) d\xi.$$

Recall that $\int \xi^{2k} \widehat{b}(\xi) = (2\pi) D_x^{2k} b(0)$.

22 April 8th, 2021

22.1 Steepest Descent

Recall for Steepest Descent, we had

$$J(\lambda) = \int \chi(x) e^{-\lambda\psi(x)} a(x) dx + O(e^{-\lambda(\psi(x_0)+a)})$$

for $\chi \in C_c^\infty$ a smooth bump function near x_0 . We then wrote $\psi(x) = \psi(x_0) + 1/2y(x)^2$, which gives a nice function $x \mapsto y(x)$ with $y(x_0) = 0$ and $y'(x_0) \neq 0$.

Applying this, we had $J(\lambda) = e^{-\lambda\psi(x_0)} \int b(y) e^{-\lambda y^2/2} dy$, $b \in C_c^\infty$ near 0 and $b(x_0) = \frac{a(x_0)}{(\psi'(x_0))^{1/2}}$.

Then, via Plancherel, we obtained

$$J(\lambda) = (2\pi)^{-1/2} \int \widehat{b}(\xi) e^{-\xi^2/2\lambda} d\xi / \sqrt{\lambda}.$$

which has a finite expansion

$$(2\pi)^{-1/2} \int \widehat{b}(\xi) \sum_{k=0}^{N-1} 1/k! (-1/2\lambda)^k \xi^{2k} d\xi / \sqrt{\lambda} + C \int \widehat{b}(\xi) \xi^{2N} e^{-1/2(\theta\xi)^2} d\xi / \sqrt{\lambda}.$$

Then, note that $\int \xi^{2k} \widehat{b}(\xi) d\xi = 2\pi D_y^{2k} b(0)$, so we obtain

$$(2\pi)^{1/2} / \sqrt{\lambda} \sum_0^{N-1} 1/k! (2\pi)^{-k} \partial_y^{2k} b(0) + C_N \lambda^{-N-1/2} \|(\widehat{D^{2N}b})\|_1.$$

Then, we estimate

$$\|(\widehat{\partial^{2N}b})\|_1 \leq c(\|\partial^{2n+2}b\|_{L^1} + \|\partial^{2N}b\|_{L^1}) \leq C \sup_{|\alpha| \leq 2N+2} |\partial^\alpha b|.$$

$$J(\lambda) = e^{-\lambda\psi(x_0)} (2\pi)^{1/2} \sum_{k=0}^{N-1} 1/k! (2\pi)^{-k} \partial_y^{2k} b(0) + O(\lambda^{-N-1/2}).$$

Putting this together, If ψ has a unique non-degenerate minimum at x_0 , $\psi'(x_0) = 0$, $\psi''(x_0) > 0$, $a \in C_0^\infty(\mathbb{R})$, we gave

$$\int a(x) e^{-\lambda\psi(x)} dx = e^{-\lambda\psi(x_0)} (2\pi/\psi''(x_0))^{1/2} 1/\sqrt{\lambda} (a(x_0) + \lambda^{-1}a_1 + \dots + \lambda^{-N-1}a_{N-1} + O(\lambda^{-N-1/2})).$$

22.2 Stirling's Formula

Recall the Gamma function

$$\Gamma(s) = \int_0^\infty e^{-t} t^s dt / t.$$

We wish to examine this as $s \rightarrow \infty$.

We rewrite this as

$$e^{-t} t^s dt / t = e^{-t+s \log t} dt / t = e^{s \log x} e^{-s(x \log x)} dx / x.$$

Then, $\Gamma(s) = s^s \int_0^\infty e^{-s(x-\log x)} \frac{dx}{x}$. The phase $\psi(x) = x - \log x$. Note that it achieves a minimum at $x = 1$ and $\psi''(x) > 0$.

Take χ to be a bump function at 1. Then,

$$\int_0^1 (1 - \chi(x))e^{-s(x - \log x)} dx/x = O(e^{-(s(1+a))}),$$

where the a is the point on the phase where χ removes the support. Similarly,

$$\int_1^\infty (1 - \chi(x))e^{-s(x - \log x)} dx/x = O(e^{-s(1+b)}).$$

We see that

$$\Gamma(s) = s^s \left[\int \chi(x) e^{-s(x - \log x)} dx/x + O(e^{-s \min(1+a, 1+b)}) \right].$$

The first term is exactly suitable for the steepest descent method.

Namely, $\psi = x - \log x$, $a(x) = 1/x$. $\psi'(1) = 0$, $\psi''(1) = 1$, $a(1) = 1$, so we obtain the estimate

$$\Gamma(s) = s^s e^{-s} [\sqrt{2\pi} s^{-1/2} + a_1 s^{-3/2} + \dots + O(s^{-N-1/2})].$$

The leading term is exactly Stirling:

$$\Gamma(s) = \sqrt{2\pi} s^{s-1/2} e^{-s} (1 + O(1/s)).$$

23 April 13th, 2021

23.1 Stationary Phase

Take $I(\lambda) = \int e^{i\lambda\varphi(x)} a(x) dx$, $\varphi \in C^\infty(\mathbb{R}; \mathbb{R})$, $a \in C_c^\infty(\mathbb{R})$.

Lemma 23.1

Suppose $|\psi'(x)| > 0$ on $\text{supp}(a)$. Then for all $K \Subset X$, N , for all $a \in C_c^\infty(K)$, there exists C such that

$$|I(\lambda)| \leq C \sup_{|\alpha| \leq N} |\partial^\alpha a| \lambda^{-N}.$$

Proof. Define ${}^tL = \frac{1}{i|\varphi'|^2} \psi' \cdot \partial_x$. Note that $1/\lambda {}^tL(e^{-\lambda\varphi}) = e^{-\lambda\varphi}$. It follows that

$$I(\lambda) = \lambda^{-N} \int e^{i\lambda\varphi} (L^N a)(x) dx \leq |\text{supp } a| \lambda^{-N} \sup |L^N a| \leq C_N \lambda^{-N} \sup_{|\alpha| \leq N} |\partial^\alpha a|.$$

□

Remark 23.2. The converse is open for higher dimensions.

This means that the contributions to the integral come from stationary points.

Theorem 23.3 (Stationary Phase(dim 1))

Suppose $X = (\alpha, \beta)$, $a \in C_c^\infty(X)$ and φ has a unique critical point in (α, β) , $\varphi'(x_0) = 0$, $\varphi''(x_0) \neq 0$. Then

$$I(\lambda) = \frac{\sqrt{2\pi}}{|\varphi''(x_0)|^{1/2}} e^{i\frac{\pi}{4} \text{sgn} \varphi''(x_0)} \lambda^{-1/2} (a_0 + \lambda^{-1} a_1 + \lambda^{-2} a_2 + \dots + \lambda^{-(N-1)} a_{N-1}) + S_N(\lambda),$$

where $a_0 = a(x_0)$ and $|S_N(\lambda)| \leq \lambda^{-1/2-N} \sup_{k \leq 2N+2} |\partial_x^k a|$.

Proof. First, note that $I(\lambda) = \int \chi(x) a(x) e^{i\lambda\varphi(x)} dx + O(\lambda^{-\infty})$, where χ is a smooth bump function near x_0 . We make a change of variables $\varphi(x) = \varphi(x_0) + \epsilon \frac{1}{2} (y(x))^2$, where $y(x_0) = 0$, $y'(x_0) = |\varphi''(x_0)|^{1/2} \neq 0$, $\epsilon = \text{sgn}(\varphi''(x_0))$. Assume $\varphi(x_0) = 0$.

Hence, $I(\lambda) = \int b(y) e^{i\lambda\epsilon y^2/2} dy + O(\lambda^{-\infty})$, where $b(y) = a(x(y)) \chi(x(y)) |dx/dy|$. Leaving out the $O(\lambda^{-\infty})$, we have $I(\lambda) = \langle e^{i\lambda\epsilon y^2/2}, b \rangle = \frac{1}{2\pi} \langle \widehat{e^{i/2\epsilon\lambda y^2}}, \widehat{b} \rangle$.

Now, we calculate the Fourier transform of $e^{i\epsilon\lambda y^2/2}$. We can write this as

$$\begin{aligned} \lim_{\delta \rightarrow 0^+} \int e^{i\epsilon/2\lambda y^2} e^{-1/2\delta y^2} e^{-iy\xi} dy &= \lim_{\delta \rightarrow 0^+} \int e^{-1/2(\delta - i\epsilon\lambda)(y + i\xi/(\delta - i\epsilon\lambda))^2 + \xi^2/(2(\delta + i\epsilon\lambda))} dy \\ &= \frac{\sqrt{2\pi}}{\sqrt{\delta - i\epsilon\lambda}} e^{-\xi^2/(2(\delta - i\epsilon\lambda))} \\ &\rightarrow \frac{\sqrt{2\pi}}{\sqrt{\lambda}} e^{i\pi/4\epsilon}. \end{aligned}$$

We obtain the Fourier Transform $\sqrt{2\pi/\lambda} e^{i\pi/4} e^{-i/2\epsilon\xi^2/\lambda}$.

Remark 23.4. There are probably sign errors above.

$$I(\lambda) = \frac{1}{2\pi} \frac{\sqrt{2\pi}}{\sqrt{\lambda}} e^{i\epsilon\pi/4} \int_{\mathbb{R}} e^{-i\epsilon\xi^2/2\lambda} \widehat{b}(\xi) d\xi.$$

Expanding the Taylor series of the exponential term, we have

$$\frac{1}{2\pi} \frac{\sqrt{2\pi}}{\sqrt{\lambda}} e^{i\epsilon\pi/4} \int_{\mathbb{R}} \sum_{k < N} \frac{1}{k!} (-i\epsilon/2\lambda)^k \xi^{2k} \widehat{b}(\xi) d\xi + \lambda^{N-1/2} \int O(\epsilon^{2N}) |\widehat{b}(\xi)| d\xi.$$

Then, note that $1/2\pi \int \xi^{2k} \widehat{b}(\xi) d\xi = (D_y^{2k} b)(0)$, so we obtain

$$\sqrt{\frac{2\pi}{\lambda}} e^{i\epsilon\pi/4} \sum_{k < N} \lambda^{-k} (-i\epsilon/2)^k D_y^{2k} b(0) + \lambda^{-N-1/2} \|\xi^{2N} \widehat{b}\|_{L^1},$$

and note that the last term is bounded by $C \sup_{k \leq 2N+2} |\partial^k b|$ (a Sobolev type estimate). This exactly gives our desired expansion. \square

23.2 Stationary Phase in Higher Dimensions

To proceed, we need an analog of $\varphi'(x_0) = 0, \varphi''(x_0) \neq 0 \Rightarrow \varphi(x) = \varphi(x_0) + \epsilon/2(y(x))^2$. For this, we use the Morse Lemma.

Theorem 23.5 (Morse Lemma)

Suppose $\varphi \in C^\infty(\mathbb{R}^n; \mathbb{R})$ and $D\varphi(x_0) = 0$ and $D^2\varphi(x_0)$ is non-degenerate. Then, there exists a transformation χ from a neighborhood of x_0 to a neighborhood of x such that $\varphi \circ \chi^{-1}(y) = \varphi(x_0) + 1/2(y_1^2 + \dots + y_r^2 - y_{r+1}^2 - \dots - y_n^2)$ where $(r, n-r)$ is the signature of $D^2\varphi(x_0)$; i. e. r is the number of positive eigenvalues of $D^2\varphi(x_0)$.

Proof. Without loss of generality, assume that $x_0 = 0, \varphi(x_0) = 0$. Then, we can diagonalize $D^2\varphi(0) = {}^t U \Lambda U$ where we order the eigenvalues $\lambda_1, \dots, \lambda_r > 0, \lambda_{r+1}, \dots, \lambda_n < 0$. Then, we can write $\varphi(x) = 1/2 \langle \varphi''(0)x, x \rangle + O(|x|^3) = 1/2 \langle \Lambda Ux, Ux \rangle + O(|Ux|^3)$, where $\tilde{x} = |\Lambda|^{1/2} Ux$. But this is just $1/2(\tilde{x}_1^2 + \dots + \tilde{x}_r^2 - \tilde{x}_{r+1}^2 - \dots - \tilde{x}_n^2) + O(|\tilde{x}|^3)$.

We write \tilde{x} as x . We can write $\varphi(x) = \int_0^1 (1-t) \partial_t^2(\varphi(tx)) dt$ (this follows from integration by parts). We can write this as $1/2 \sum q_{jk}(x) x_j x_k$ where $q_{jk}(x) = 2 \int_0^1 (1-t) \frac{\partial^2 \varphi}{\partial x_j \partial x_k}(tx) dt, q_{jk} = q_{kj}$.

Then, $q_{jk}(0) = \frac{\partial^2 \varphi}{\partial x_j \partial x_k}(0)$ so defining a matrix $Q(x) = (q_{jk}(x))$, we have $\varphi(x) = 1/2 \langle Q(x)x, x \rangle, Q(0) = D^2\varphi(x_0)$.

We would like to find $k(x) = A(x)x$ where $A(0) = I, x \mapsto A(x) \in C^\infty$. Then, $\langle x, Q(x)x \rangle = \langle A(x)x, Q(0)A(x)x \rangle = \langle K(x), Q(0)K(x) \rangle = \langle y, Q(0)y \rangle$ where $y = K(x)$.

This is the same as $\langle x, Q(x)x \rangle = \langle x, {}^t A(x)Q(0)A(x)x \rangle$. So, we need $Q(x) = A^T(x)Q(0)A(x)$ with $A(0) = I$. To handle this, we use the implicit function theorem.

Theorem 23.6 (Implicit Function Theorem)

Given a function $F : \mathbb{R}^M \rightarrow \mathbb{R}^N$ and $F(X_0) = Y_0 \in \mathbb{R}^N, X_0 \in \mathbb{R}^M$. We solve $F(X(Y)) = Y, X(Y_0) = X_0$, we need $dF(X_0) : \mathbb{R}^M \rightarrow \mathbb{R}^N$ to be surjective (or F is a smooth submersion).

We will use $F : M(\mathbb{R}, n \times n) \rightarrow \text{Sym}(\mathbb{R}, n)$ where $F(A) = A^T Q(0)A$. Then, $F(A+H) = F(A) + dF_A(H) + o(\|H\|)$. We have $A^T Q(0)A + H^T Q(0)A + A^T Q(0)H + H^T Q(0)H$. \square

24 April 18th, 2021

24.1 Morse Lemma

Recall the Morse Lemma:

Theorem 24.1 (Morse Lemma)

Suppose $\varphi \in C^\infty(\mathbb{R}^n; \mathbb{R})$ and $D\varphi(x_0) = 0$ and $D^2\varphi(x_0)$ is non-degenerate. Then, there exists a transformation χ from a neighborhood of x_0 to a neighborhood of x such that $\varphi \circ \chi^{-1}(y) = \varphi(x_0) + 1/2(y_1^2 + \cdots + y_r^2 - y_{r+1}^2 - \cdots - y_n^2)$ where $(r, n-r)$ is the signature of $D^2\varphi(x_0)$; i. e. r is the number of positive eigenvalues of $D^2\varphi(x_0)$.

Proof. Last time, we reduced the problem to finding $x \mapsto A(x)$, where $A(x) \in \mathbb{R}^{n \times n}$ and $A(0) = I_n$. Then, if we take $Q(x) = A^T(x)Q(0)A(x)$, it follows that $x \mapsto Q(x) \in \text{Symm}(n)$, the group of symmetric real-valued matrices, where $Q(0) = Q$ is non-degenerate.

To do this, define $F : \mathbb{R}^{n \times n} \rightarrow \text{Symm}(n)$ with $F(A) = A^TQA$. We wish to find a right inverse H such that $G(Q) = I$ and $F \circ G = I_n$. Then, $A(x) = G(Q(x))$ would satisfy our conditions.

Note that $F(A + H) = F(A) + dF_A(H) + o(\|H\|^2)$. Namely,

$$(A + H)^TQ(A + H) = A^TQA + A^TQH + H^TQA + H^TQH.$$

The first term is $F(A)$, the next two give $dF_A(H)$ and the last is $o(\|H\|^2)$. Now, $dF_A : \mathbb{R}^{n \times n} \rightarrow \text{Symm}(n)$ is given by $H \mapsto QH + H^TQ = S$, note that we can find a right inverse $Q^{-1}S/2$. It follows that we can apply the implicit function theorem. \square

24.2 Higher Dimensional Method of Stationary Phase

As before, take $I(\lambda) = \int e^{i\lambda\varphi(x)}a(x)dx$ for $a \in C_c^\infty(X)$, where we assume φ has a non-degenerate critical point. We can write this as

$$\begin{aligned} &= \int e^{i\lambda\varphi(x)}\chi(x)a(x)dx + O(\lambda^{-\infty}) \quad \text{supp } \chi \in B_\epsilon(x_0) \\ &= \int e^{-i\lambda/2\langle Qx, x \rangle}b(x)dx + O(\lambda^{-\infty}), \quad b(x) = a(K^{-1}(x)) \left| \frac{dK^{-1}}{dx} \right| \end{aligned}$$

by applying the Morse Lemma. If we take the Fourier Transform of the Gaussian and apply the Blancherel Formula, we obtain

$$\begin{aligned} &= \frac{\lambda^{-n/2}(2\pi)^{n/2}}{(2\pi)^n |\det Q|^{1/2}} e^{i\pi/4 \text{sgn } Q} \int \sum_{k < N} 1/k! (\langle Q^{-1}\xi, \xi \rangle / (2\lambda i))^{k/2} \widehat{b}(\xi) + \lambda^{-n/2-N} \int O(\langle \xi \rangle^{2N}) |\widehat{b}(\xi)| d\xi \\ &= \frac{\lambda^{-n/2}(2\pi)^{n/2}}{(2\pi)^n |\det Q|^{1/2}} e^{i\pi/4 \text{sgn } Q} \sum_{k < N} \frac{1}{k!} \left[\left(\frac{\langle Q^{-1}D_x, D_x \rangle}{2\lambda i} \right)^k b \right] (0) + \lambda^{-N-n/2} S_n(\lambda), \end{aligned}$$

where

$$|S_N(\lambda)| \leq C \|\langle \xi \rangle^{2n} \widehat{b}\|_q \leq C \sum_{|\alpha| \leq 2N+n+1} \|\partial^\alpha b\|_\infty.$$

Example 24.2

Take $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$, $Q = \begin{pmatrix} 0 & -I \\ -I & 0 \end{pmatrix}$. Note that $Q^{-1} = Q$. Then, $1/2 \langle Q(x, y), (x, y) \rangle =$

$-x \cdot y = w^2 - v^2$ where $w = \frac{x+y}{\sqrt{2}}$, $v = \frac{x-y}{\sqrt{2}}$. This corresponds to a matrix $\begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}$, which has a signature of 0. This is the nicest case for the stationary phase method.

Then, $\frac{1}{2i} \langle Q^{-1} D_{x,y}, D_{x,y} \rangle = \frac{1}{i} \sum_{i=1}^n D_{x_j} D_{y_j}$ which gives $(\lambda/2\pi)^n \int e^{-\lambda x \cdot y} u(x, y) dx dy$ for $u \in C_c^\infty(\mathbb{R}^n \times \mathbb{R}^n)$. From the method of stationary phase, we have

$$\sum_0^{N-1} \frac{1}{k!} \frac{1}{\lambda^k} \left(\sum 1/i D_{x_j} D_{y_j} \right)^k u(0, 0) + S_n \lambda^{-N},$$

$$\|S_N\| \leq C \sum_{|\alpha+\beta| \leq 2n+1} \|\partial_x^\alpha \partial_y^\beta (\partial_x \cdot \partial_y)\|_\infty$$

24.3 General Schrodinger Operators

We find a fundamental solution to

$$(i\partial_t + \frac{1}{2} \langle Q^{-1} D_x, D_x \rangle) E = \delta_0(t) \delta_0(x).$$

For $Q = -2I$, we have the free Schrodinger Equation.

We require Q to be a symmetric, non-degenerate matrix. If we take a spacial fourier transform, we obtain

$$\delta_0(t) = (i\partial_t + \langle Q^{-1} \xi, \xi \rangle / 2) \widehat{E}_x = i\partial_t (e^{-it/2 \langle Q^{-1} \xi, \xi \rangle} \widehat{E}),$$

a first-order ODE. Solving this, we obtain

$$\widehat{E}_x(t, \xi) = e^{it/2 \langle Q^{-1} \xi, \xi \rangle} H(t) / i,$$

where H is the heaviside function. Finally, if we take an inverse spacial fourier transform we obtain

$$E = \frac{|\det Q|^{1/2}}{i(2t\pi)^{n/2}} e^{i\pi/4 \operatorname{sgn} Q} e^{i \langle Qx, x \rangle / (2it)}.$$

25 April 20th, 2021

25.1 Pseudodifferential Operators

Definition 25.1. An operator A is a pseudodifferential operator of order $\leq m$, $A \in \Psi_{\rho,\delta}^m(X)$ ($\rho > 0$, $\delta < 1$), if there exists $a \in S_{\rho,\delta}^m(X \times X \times \mathbb{R}^n)$ such that for $u \in C_c^\infty(X)$,

$$Au(x) = (2\pi)^{-n} \iint e^{i(x-y)\xi} a(x, y, \xi) u(y) dy d\xi$$

as an oscillatory integral.

- If we take $A = \sum_{|\alpha| \leq m} a_\alpha D_x^\alpha$, then $a(x, y, \xi) = \sum_{|\alpha| \leq m} a_\alpha(x) \xi^\alpha$ is the corresponding symbol.
- $(I - \Delta) : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ has an inverse $(I - \Delta)^{-1} : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n) \in \Psi_{1,0}^{-2}$, by taking $a(x, y, \xi) = (1 + |\xi|^2)^{-1}$.
- $P(Eu) = u$, $u \in \mathcal{E}'(\mathbb{R}^{n+1})$, with $P = \partial_t - \Delta + 1$. Then, $E \in \Psi_{1/2,0}^{-1}(\mathbb{R}^{n+1})$ with the (dreaded) HW symbol $a = (i\tau + |\xi|^2 + 1)^{-1}$.
- $(i\partial_t + \Delta)(Eu) = u$, $u \in \mathcal{E}'(\mathbb{R}^{n+1})$. We showed that $a = e^{it|\xi|^2} H(t)$, which is not in any symbol class.

Recall from before,

$$Au(y) = \int K_a(x, y) u(x) dx, \quad K_a \in \mathcal{D}'(X \times X), u \in C_c^\infty(X).$$

Then, $Au \in \mathcal{D}'(X)$, $\langle Au, v \rangle = \langle K_a, u \otimes v \rangle$, $v \in C_c^\infty(X)$. If $K_A = I(a, \varphi)$, then $(x, \theta) \mapsto \varphi(x, y, \theta)$ takes $C_c^\infty(X) \rightarrow C^\infty(X)$ and $(y, \theta) \mapsto \varphi(x, y, \theta)$ takes $\mathcal{E}'(X) \rightarrow \mathcal{D}'(X)$.

Recall the singular support of K_A is contained in the set $\{(x, y) : \exists \delta, \varphi_\delta(x, y, \delta) = 0\}$. Note that if $A \in \Psi_{\rho,\delta}^m$, then $A : C_c^\infty(X) \rightarrow C^\infty(X)$ and $A : \mathcal{E}'(X) \rightarrow \mathcal{D}'(X)$. In this case, the singular support is contained in $\Delta(X \times X) = \{(x, x) : x \in X\}$.

Proposition 25.2 (Semilocality)

If $A : \mathcal{E}'(X) \rightarrow \mathcal{D}'(X)$, then $\text{sing supp } K_A \subset \text{sing supp } u$.

Proof. Take $x_0 \in X \setminus \text{sing supp } u$. Choose $\varphi \in C_c^\infty(X)$ with $\varphi \equiv 1$ near x_0 , $\psi \equiv 1$ near $\text{sing supp } u$ and $\text{supp } \varphi \cap \text{supp } \psi = \emptyset$. Then, $Au = A(1 - \psi)u + A\psi u = A\psi u + g$ with $g \in C^\infty(X)$. It follows that $\varphi Au = \varphi A\psi u + \varphi g$, $\varphi g \in C_c^\infty(X)$. Since φ is 1 near x_0 , it suffices to show that $\varphi Au \in C^\infty$, since it would follow x_0 cannot be in the singular support of Au .

Define $K_{\varphi A \psi}(x, y) = \varphi(x) K_A(x, y) \psi(y)$. Since $\text{supp } K_{\varphi A \psi} \cap \Delta(X \times X) = \emptyset$, we have $K_{\varphi A \psi} \in C_c^\infty$, which implies that $\varphi A\psi u \in C^\infty$, which implies that $\varphi Au \in C^\infty$. \square

Definition 25.3. A is a smoothing operator if A takes $\mathcal{E}'(X) \rightarrow C^\infty(Y)$ and $C_c^\infty(Y) \rightarrow \mathcal{D}'(X)$, or equivalently $K_A \in C^\infty(X \times Y)$. We define $\Psi^{-\infty}$ to be the class of smooth operators.

Proposition 25.4

$A \in \Psi^{-\infty}(X)$ if and only if $a \in S^{-\infty}(X \times X \times \mathbb{R}^n)$ satisfying the pseudodifferential operator definition.

Proof. For the forward direction, $K_A(x, y) = C \int e^{i(x-y)\xi} a(x, y, \xi) d\xi \in C^\infty(X \times X)$. \square